arXiv:math/0609715v1 [math.QA] 26 Sep 2006

geometry and reducibility of induction of Hopf group co-algebra

A.S.Hegazi[1] F.Ismail[2] and M.M. Elsofy[3]

[1] Department of Mathematics, Faculty of Science, Mansoura University, Egypt.
Email: Hegazi@mans.edu.eg

[2] Department of Mathematics, Faculty of Science, Cairo University, Egypt.
Email: fatma-ism@hotmail.com

[3] Department of Mathematics, Faculty of Science, Fayoum University, Egypt.
Email: melsoffyy@yahoo.com

Abstract

In this work we study the induction (induced and coinduced)theory for Hopf group coalgebra. We define a substructure B of a Hopf group coalgebra H, called subHopf group coalgebra. Also, we have introduced the definition of Hopf group suboalgebra and group coisotropic quantum subgroup of H. The properties of the algebraic structure of the induced and coinduced are given. Moreover, a framework of the geometric interperation and simplicity theory of such representation strructure are stuided.

1 Introduction

The induced representation of quantum group (quasitriangular Hopf algebra [M] and [Mon]) is introduced by Gonzalez-Ruiz, L. A. Ibort [G-I] and Ciccoli [C]. Recently, Hegazi, Agawany, F. Ismail and I. Saleh in [H], have succeeded to give a new algebric structure for quantum subgroups and subquantum groups, that is a subspase B of a bialgebra H is called a sub-bi-algebra if the restriction to B of the structure of H turns B into bialgebra. But a bi-sub-algebra of H is a different thing. It is pair (B, π) consisting of another bialgebra and surjective homomorphism $\pi : H \to B$. A relation between these two notions are given if the bialgebra allows a decomposition of the form $H = B \oplus I$ with I a bi-ideal in H. The projection π into B yields a bi-sub-algebra (B, π) in the above sense. Conversely if $B \subseteq H$ is both a sub-bi-algebra and a bi-sub-algebra of H such that $\pi^2 = \pi$, then $H = B + Ker$ π and we have a decomposition of the form $H = B \oplus I$. These construction made us able to introduce the induced representation of Hopf algebra in each case. That is a B-comodule for a sub-Hopf algebra B of a Hopf algebra H gives rise to an induced H-Hopf module and that a representation of a quantum subgroup (B, π) of a quantum group H induced a Hopf representation of H. This procedure realizes a quantum group induced representation.

Recently, Turaev [T]and Virelizier [V-V1] give use a new definition for a generalization of Hopf algebra, for Hopf algebra structure see [S] and [Mon], Hopf group coalgebra. These generalization gives us quantum group structure this structure is of great importance in Homotopy theory in quantum field theory [T].

In this paper, unless otherwise, every thing takes place over a field K, and K-space means vector space over K. A map f from a space V into a space W always means linear map over K. The tensor product $V \otimes W$ is understood to be $V \otimes_K W$, $I : V \rightarrow V$ always denotes the identity map, and the transposition map $\tau : V \otimes W \rightarrow W \otimes V$ is defined by $\tau(v \otimes w) = w \otimes v$ for $v \in V, w \in W$. Let $f : C \rightarrow D$ be a map. Then $f^{\ *} : D^* \rightarrow C^*$ is a map, where $f^*(\phi)(c) = \phi(f(c))$ for all $\phi \in D^*, c \in C$.

Recently, Virelizier [V-V1] studied the algebraic properties of the Hopf $\pi-$coalgebra. Now, let us give some basic definitions about Hopf $\pi-$coalgebra. For group π, a π-coalgebra is a family $C = \{C_\alpha\}_{\alpha \in \pi}$ of K-spaces endowed with a family K-maps (comultiplication) $\Delta = \{\Delta_{\alpha,\beta} : C_{\alpha\beta} \rightarrow C_\alpha \otimes C_\beta\}_{\alpha,\beta\epsilon\pi}$, K-map (counit) $\varepsilon : C_1 \rightarrow k$ such that the following diagrams are commute:

$$\begin{tikzcd}
C_{\alpha\beta\gamma} \arrow[r, "\Delta_{\alpha,\beta\gamma}"] \arrow[d, "\Delta_{\alpha\beta,\gamma}"'] & C_\alpha \otimes C_{\beta\gamma} \arrow[d, "I_\alpha \otimes \Delta_{\beta,\gamma}"] \\
C_{\alpha\beta} \otimes C_\gamma \arrow[r, "\Delta_{\alpha,\beta} \otimes I_\gamma"'] & C_\alpha \otimes C_\beta \otimes C_\gamma
\end{tikzcd}$$

$$(\Delta_{\alpha,\beta} \otimes I_\gamma\ )\Delta_{\alpha\beta,\gamma} = (I_\alpha \otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma}$$

$$\begin{tikzcd}
& C_1 \otimes C_\alpha & & C_\alpha \otimes C_1 \arrow[dr, "I_\alpha \otimes \epsilon"] & \\
K \otimes C_\alpha & & & & C_\alpha \otimes K \\
& C_\alpha \arrow[u, "\Delta_{1,\alpha}"'] \arrow[ul, "\sim_\alpha"] & & C_\alpha \arrow[u, "\Delta_{\alpha,1}"] \arrow[ur, "\sim_\alpha"'] &
\arrow[from=1-2, to=2-1, "\epsilon \otimes I_\alpha"']
\end{tikzcd}.$$

$$(\epsilon \otimes I_\alpha)\Delta_{1,\alpha} = \sim_\alpha = (I_\alpha \otimes \epsilon)\Delta_{\alpha,1}$$

A Hopf $\pi-$coalgebra is a π-coalgebra $(H = \{H_\alpha\}_{\alpha\in\pi}, \Delta, \epsilon)$ with a family $S = \{S_\alpha : H_\alpha \to H_{\alpha^{-1}}\}_{\alpha\in\pi}$ of K-maps such that

1. H_α is an algebra with multiplication μ_α and unit $\eta_\alpha(1_K)$ for all $\alpha \in \pi$;
2. $\Delta_{\alpha,\beta}$, ϵ are algebra maps for all $\alpha, \beta \in \pi$,
3. The antipode S satisfy
   $\mu_\alpha(S_{\alpha^{-1}} \otimes I_\alpha)\Delta_{\alpha^{-1},\alpha} = 1_\alpha\epsilon = \mu_\alpha(I_\alpha \otimes S_{\alpha^{-1}})\Delta_{\alpha,\alpha^{-1}}$

A Hopf π-coalgebra H is of finite type when every H_α is finite dimensional. Note that it does not means that $\oplus_{\alpha\in\pi}H_\alpha$ is finite dimensional (unless $H_\alpha = 0$ for all but a finite number of $\alpha \in \pi$). The notion of Hopf $\pi-$coalgebra is not self dual. Note, $(H_1, \Delta_{1,1}, \epsilon)$ is a (classical) Hopf algebra.

Let C be a π-coalgebra. A right π-comodule over C is a family $M = \{M_\alpha\}_{\alpha\in\pi}$ of K-spaces endowed with a family $\theta = \{\theta_{\alpha,\beta} : M_{\alpha\beta} \to M_\alpha \otimes C_\beta\}_{\alpha,\beta\epsilon\pi}$ of K-maps such that the following diagrams are commute:

$$\begin{tikzcd}
M_{\alpha\beta\gamma} \arrow[r, "\theta_{\alpha,\beta\gamma}"] \arrow[d, "\theta_{\alpha\beta,\gamma}"'] & M_\alpha \otimes C_{\beta\gamma} \arrow[d, "I_\alpha \otimes \Delta_{\beta,\gamma}"] \\
M_{\alpha\beta} \otimes C_\gamma \arrow[r, "\theta_{\alpha,\beta}\otimes I_\gamma"] & C_\alpha \otimes C_\beta \otimes C\gamma
\end{tikzcd}
\qquad
\begin{tikzcd}
M_\alpha \arrow[dr, "\theta_{\alpha,1}"] \arrow[dd, "\sim_{M_\alpha}"'] & \\
& M_\alpha \otimes C_1 \arrow[dl, "I_\alpha \otimes \epsilon"] \\
M_\alpha \otimes K &
\end{tikzcd}$$

$$(\theta_{\alpha,\beta} \otimes I_\gamma\ )\theta_{\alpha\beta,\gamma} = (I_\alpha \otimes \Delta_{\beta,\gamma})\theta_{\alpha,\beta\gamma} \qquad (I_\alpha \otimes \epsilon)\theta_{\alpha,1} = \sim_{M_\alpha}$$

A right Hopf π-comodule over H is a right π-comodule $M = \{M_\alpha\}_{\alpha\in\pi}$ by coaction $\theta = \{\theta_{\alpha,\beta}\}_{\alpha,\beta\epsilon\pi}$ such that

1. M_α is $H_\alpha-$module by action ρ_α for all $\alpha \in \pi$
2. The following diagram, for all $\alpha, \beta, \gamma \in \pi$, is commute:

$$\begin{tikzcd}
M_{\alpha\beta} \otimes H_{\alpha\beta} \arrow[r, "\rho_{\alpha\beta}"] \arrow[d, "\theta_{\alpha,\beta} \otimes \Delta_{\alpha,\beta}"'] & M_{\alpha\beta} \arrow[r, "\theta_{\alpha,\beta}"] & M_\alpha \otimes H_\beta \\
(M_\alpha \otimes H_\beta) \otimes (H_\alpha \otimes H_\beta) \arrow[rr, "I_\alpha\otimes\tau\otimes I_\beta"] & & (M_\alpha \otimes H_\alpha) \otimes (H_\beta \otimes H_\beta) \arrow[u, "\rho_\alpha \otimes \mu_\beta"']
\end{tikzcd}$$

i.e., $\theta_{\alpha,\beta}\ \rho_{\alpha\beta} = (\rho_\alpha \otimes \mu_\beta)(I_\alpha \otimes \tau \otimes I_\beta)(\theta_{\alpha,\beta} \otimes \Delta_{\alpha,\beta})$

Let C be a $\pi-$coalgebra. A $\pi-$coideal of C is a family $V = \{V_\alpha\}_{\alpha\in\pi}$ such that

1. V_α is subspace of C_α for all $\alpha \in \pi$,
2. $\Delta_{\alpha,\beta}(V_{\alpha\beta}) \subset V_\alpha \otimes C_\beta + C_\alpha \otimes V_\beta$ for $\forall\alpha,\beta \in \pi$. and $\epsilon_1(V_1) = 0$.

A Hopf π-coideal of H is a family of K-spaces $V = \{V_\alpha\}_{\alpha\in\pi}$ such that V is π-coideal of H, V_α is ideal of H_α and $S_\alpha(V_\alpha) \subseteq V_{\alpha^{-1}}$ for all $\alpha \in \pi$.

A family $A = \{A_\alpha\}_{\alpha\in\pi}$ is called subHopf $\pi-$coalgebra of H if (A, Δ, ϵ) is $\pi-$coalgebra of H, A_α is a subalgebra of H_α and $S_\alpha(A_\alpha) \subseteq A_{\alpha^{-1}}$ $\forall\alpha \in \pi$. A family $A = \{A_\alpha\}_{\alpha\in\pi}$ of a subalgebra of Hopf π-coalgebra of H is called isolated if there exist a family $I = \{I_\alpha\}_{\alpha\in\pi}$ of Hopf π- coideal of H such that $H_\alpha = A_\alpha \oplus I_\alpha$ $\forall\alpha \in \pi$.

We will call Hopf $\pi-$subcoalgebra of H any pair (C, σ) such that $C = \{C_\alpha\}_{\alpha\in\pi}$ is a Hopf $\pi-$coalgebra and $\sigma = \{\sigma_\alpha : H_\alpha \to C_\alpha\}_{\alpha\in\pi}$ family of algebra epimorphisms which satisfies for $\alpha, \beta \in \pi$

1. $\Delta^C_{\alpha,\beta}\sigma_{\alpha\beta} = (\sigma_\alpha \otimes \sigma_\beta)\Delta^H_{\alpha,\beta}$, i.e., the following diagram is commute

$$\begin{tikzcd}
H_{\alpha\beta} \arrow[r, "\sigma_{\alpha,\beta}"] \arrow[d, "\Delta^H_{\alpha,\beta}"'] & C_{\alpha\beta} \arrow[d, "\Delta^C_{\alpha,\beta}"] \\
H_\alpha \otimes H_\beta \arrow[r, "\sigma_\alpha \otimes \sigma_\beta"'] & C_\alpha \otimes C_\beta
\end{tikzcd}$$

2. $\varepsilon^C\sigma_1 = \varepsilon^H$, i.e., the following diagram is commute

$$\begin{tikzcd}
H_1 \arrow[r, "\sigma_1"] \arrow[dr, "\epsilon^H"'] & C_1 \arrow[d, "\epsilon^C"] \\
 & K
\end{tikzcd}$$

3. $S^C_\alpha\sigma_\alpha = \sigma_{\alpha^{-1}}S^H_\alpha$ i.e., the following diagram is commute

$$\begin{tikzcd}
H_\alpha \arrow[r, "\sigma_\alpha"] \arrow[d, "S^H_\alpha"'] & C_\alpha \arrow[d, "S^C_\alpha"] \\
H_{\alpha^{-1}} \arrow[r, "\sigma_{\alpha^{-1}}"'] & C_{\alpha^{-1}}
\end{tikzcd}$$

A pair (C, σ) is called left $\pi-$coisotropic quantum subgroup of H if

1. C is $\pi-$coalgebra,
2. C_α is left H_α-module by ω_α for all $\alpha \in \pi$,
3. $\sigma = \{\sigma_\alpha : H_\alpha \to C_\alpha\ \}_{\alpha\in\pi}$ family of surjective linear maps such that
   (a) σ_α is left H_α-module map for all $\alpha \in \pi$,
   (b) $\Delta^C_{\alpha,\beta}\sigma_{\alpha\beta} = (\sigma_\alpha \otimes \sigma_\beta)\Delta^H_{\alpha,\beta}$ and $\varepsilon^C \circ \sigma_1 = \varepsilon^H$.

If $g = \{g_\alpha : C_\alpha \to H_\alpha\}_{\alpha\in\pi}$, is a family of linear maps, its convolution inverse is a family of linear maps $g^{-1} = \{g^{-1}_\alpha : C_{\alpha^{-1}} \to H_\alpha\}_{\alpha\in\pi}$ such that

$$\mu_\alpha(g_\alpha \otimes g^{-1}_\alpha)\Delta^C_{\alpha,\alpha^{-1}}(c) = \epsilon^C(c)1_{H_\alpha} = \mu_\alpha(g^{-1}_\alpha \otimes g_\alpha)\Delta^C_{\alpha^{-1},\alpha}(c).$$

Proposition 1.1. *Every isolated subHopf $\pi-$coalgebra is $\pi-$coisotropic quantum subgroup.*

Proof. Clear from definition. ∎

A left $\pi-$coisotropic quantum subgroup (C, σ) of H is said to have a left section if there exist a family of linear, convolution invertible, maps $g = \{g_\alpha : C_\alpha \to H_\alpha\}_{\alpha\in\pi}$ such that

1. $g_\alpha(\sigma_\alpha(1)) = 1$,
2. for $\alpha \in \pi, u \in H_1, c \in C_\alpha$ and $v \in \sigma^{-1}_\alpha(c)$ we have
   $(\sigma_1 \otimes I^H_\alpha)(\mu_1 \otimes I_\alpha)(I_1 \otimes \tau)(\Delta^H_{1,\alpha} g_\alpha \otimes I_1)(c \otimes u)$
   $= (I_1 \otimes g_\alpha)(\sigma_1 \otimes \sigma_\alpha)(\mu_1 \otimes I_\alpha)(I_1 \otimes \tau)(\Delta^H_{1,\alpha} \otimes I_1)(v \otimes u)$
3. for $\alpha \in \pi, u \in H_1, c \in C_{\alpha^{-1}}$ and $v \in \sigma^{-1}_{\alpha^{-1}}(c)$ we have
   $(\sigma_1 \otimes I^H_\alpha)(\mu_1 \otimes I_\alpha)(I_1 \otimes \tau)(\Delta^H_{1,\alpha} g^{-1}_\alpha \otimes I_1)(c \otimes u)$
   $= (I_1 \otimes g^{-1}_\alpha)(\sigma_1 \otimes \sigma_{\alpha^{-1}})(\mu_1(S^H_1 \otimes I_1) \otimes I_{\alpha^{-1}})(I_1 \otimes \tau)(\tau\Delta^H_{\alpha^{-1},1} \otimes I_1)(v \otimes u)$

Theorem 1.2. *Let C be $\pi-$coalgebra, V be a $\pi-$coideal and $\sigma = \{\sigma_\alpha : C_\alpha \longrightarrow E_\alpha = C_\alpha/V_\alpha\}_{\alpha\in\pi}$ be a family of the natural linear maps onto the quotient vector spaces. Then $E = \{E_\alpha = C_\alpha/V_\alpha\}_{\alpha\in\pi}$ has a unique $\pi-$coalgebra structure such that σ is a $\pi-$coalgebra maps.*

Proof. Since V is $\pi-$coideal, $\epsilon_1(V_1) = 0$ implies $V_1 \subseteq Ker\epsilon_1$. Hence there exist unique linear map $\overline{\epsilon}_1$ making the following diagram commutes

$$\begin{tikzcd}
C_1 \arrow[rr, "\epsilon_1"] \arrow[dr, "\sigma_1"'] & & K \\
& E_1 \arrow[ur, "\overline{\epsilon_1}"'] &
\end{tikzcd}$$

Let

$$T = \{T_{\alpha,\beta} : C_{\alpha\beta} \xrightarrow{\Delta_{\alpha,\beta}} C_\alpha \otimes C_\beta \xrightarrow{\sigma_\alpha\otimes\sigma_\beta} E_\alpha \otimes E_\beta\}.$$

Since $Ker\sigma_\alpha = V_\alpha$, then $Ker(\sigma_\alpha \otimes \sigma_\beta) = C_\alpha \otimes V_\beta + V_\alpha \otimes C_\beta \supseteq \Delta_{\alpha,\beta}(V_{\alpha\beta})$, so $V_{\alpha\beta} \subseteq KerT_{\alpha,\beta}$ and since $T_{\alpha,\beta} : C_{\alpha\beta} \longrightarrow E_\alpha \otimes E_\beta$ then there exist unique linear maps $\overline{\Delta}_{\alpha,\beta} : C_{\alpha\beta}/V_{\alpha\beta} = E_{\alpha\beta} \longrightarrow E_\alpha \otimes E_\beta$ such that the following diagram commutes

$$\begin{tikzcd}
C_{\alpha\beta} \arrow[rr, "T_{\alpha,\beta}"] \arrow[dr, "\sigma_{\alpha\beta}"'] & & E_\alpha \otimes E_\beta \\
& E_{\alpha\beta} \arrow[ur, "\overline{\Delta}_{\alpha,\beta}"'] &
\end{tikzcd}$$

i.e., $\overline{\Delta}_{\alpha,\beta}\sigma_{\alpha\beta} = T_{\alpha,\beta} = (\sigma_\alpha \otimes \sigma_\beta)\Delta_{\alpha,\beta}$ for all $\alpha, \beta \in \pi$. Thus $\overline{\Delta}_{\alpha,\beta\gamma}\sigma_{\alpha\beta\gamma} = (\sigma_\alpha \otimes \sigma_{\beta\gamma})\Delta_{\alpha,\beta\gamma}$ for all $\alpha, \beta, \gamma \in \pi$. Now,

$$\begin{aligned}
&(I_\alpha \otimes \overline{\Delta}_{\beta,\gamma})\overline{\Delta}_{\alpha,\beta\gamma}\sigma_{\alpha\beta\gamma} \\
&= (I_\alpha \otimes \overline{\Delta}_{\beta,\gamma})(\sigma_\alpha \otimes \sigma_{\beta\gamma})\Delta_{\alpha,\beta\gamma} \\
&= (\sigma_\alpha \otimes \overline{\Delta}_{\beta,\gamma}\sigma_{\beta\gamma})\Delta_{\alpha,\beta\gamma} \\
&= (\sigma_\alpha \otimes (\sigma_\beta \otimes \sigma_\gamma)\Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma} \\
&= (\sigma_\alpha \otimes \sigma_\beta \otimes \sigma_\gamma)(I_\alpha \otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma} \\
&= (\sigma_\alpha \otimes \sigma_\beta \otimes \sigma_\gamma)(\Delta_{\alpha,\beta} \otimes I_\gamma)\Delta_{\alpha\beta,\gamma} \\
&= ((\sigma_\alpha \otimes \sigma_\beta)\Delta_{\alpha,\beta} \otimes \sigma_\gamma)\Delta_{\alpha\beta,\gamma} \\
&= (\overline{\Delta}_{\alpha,\beta}\sigma_{\alpha\beta} \otimes \sigma_\gamma)\Delta_{\alpha\beta,\gamma} \\
&= (\overline{\Delta}_{\alpha,\beta} \otimes I_\gamma)(\sigma_{\alpha\beta} \otimes \sigma_\gamma)\Delta_{\alpha\beta,\gamma} \\
&= (\overline{\Delta}_{\alpha,\beta} \otimes I_\gamma)\overline{\Delta}_{\alpha\beta,\gamma}\sigma_{\alpha\beta\gamma}
\end{aligned}$$

since σ_α is surjective for all $\alpha \in \pi$ then $(I_\alpha \otimes \overline{\Delta}_{\beta,\gamma})\overline{\Delta}_{\alpha,\beta\gamma} = (\overline{\Delta}_{\alpha,\beta} \otimes I_\gamma)\overline{\Delta}_{\alpha\beta,\gamma}$. Also, since $\overline{\Delta}_{\alpha,\beta}\sigma_{\alpha\beta} = (\sigma_\alpha \otimes \sigma_\beta)\Delta_{\alpha,\beta}$ implies $\overline{\Delta}_{\alpha,1}\sigma_{\alpha 1} = (\sigma_\alpha \otimes \sigma_1)\Delta_{\alpha,1}$ and so

$$\begin{aligned}
&(I_{E_\alpha} \otimes \overline{\epsilon}_1)\overline{\Delta}_{\alpha,1}\sigma_{\alpha 1} \\
&= (I_{E_\alpha} \otimes \overline{\epsilon}_1)(\sigma_\alpha \otimes \sigma_1)\Delta_{\alpha,1} \\
&= (\sigma_\alpha \otimes I_K)\underbrace{(I_{H_\alpha} \otimes \epsilon_1)\Delta_{\alpha,1}}_{} \\
&= \sigma_{\alpha 1},
\end{aligned}$$

i.e., $(I_{E_\alpha} \otimes \overline{\epsilon}_1)\overline{\Delta}_{\alpha,1}\sigma_{\alpha 1} = \sigma_{\alpha 1}$. Since σ_α is surjective for all $\alpha \in \pi$, then $(I_{E_\alpha} \otimes \overline{\epsilon}_1)\overline{\Delta}_{\alpha,1} = I_{E_\alpha}$. Similar, $(\overline{\epsilon}_1 \otimes I_{E_\alpha})\overline{\Delta}_{1,\alpha} = I_{E_\alpha}$. Thus $(E, \overline{\Delta}, \overline{\epsilon}_1)$ is $\pi-$coalgebra.

∎

2 Induced representations of Hopf group coalgebra

In this section, we study the induced representation for Hopf group coalgebra. To reach this goal we use the definitions of subHopf group coalgebra, Hopf group subcoalgebra and group coisotropic quantum subgroup.

Theorem 2.1. *Let H be a Hopf $\pi-$coalgebra and (C, σ) be a Hopf $\pi-$subcoalgebra of H. Then (C, σ) is a left $\pi-$coisotropic quantum subgroup of H.*

Proof. We define $\omega = \{\omega_\alpha = \mu_\alpha^C(\sigma_\alpha \otimes I_\alpha) : H_\alpha \otimes C_\alpha \to C_\alpha\}_{\alpha\in\pi}$ we'll prove that C_α is a left $H_\alpha-$module by ω_α. i.e., the following diagrams are commute

$$\begin{tikzcd}
H_\alpha \otimes H_\alpha \otimes C_\alpha \arrow[r, "I_\alpha \otimes \omega_\alpha"] \arrow[d, "\mu_\alpha \otimes I_\alpha"'] & H_\alpha \otimes C_\alpha \arrow[d, "\omega_\alpha"] \\
H_\alpha \otimes C_\alpha \arrow[r, "\omega_\alpha"'] & C_\alpha
\end{tikzcd}
\qquad
\begin{tikzcd}
K \otimes C_\alpha \arrow[rr, "\eta_\alpha \otimes I_\alpha"] \arrow[dr, "\sim"'] & & H_\alpha \otimes C_\alpha \arrow[dl, "\omega_\alpha"] \\
& C_\alpha &
\end{tikzcd}$$

$$\begin{aligned}
&\omega_\alpha(I_\alpha \otimes \omega_\alpha)(h \otimes k \otimes b) \\
&= \omega_\alpha(h \otimes \sigma_\alpha(k)b) \\
&= \sigma_\alpha(h)(\sigma_\alpha(k)b) \\
&= \sigma_\alpha(hk)b \\
&= \omega_\alpha(hk \otimes b) \\
&= \omega_\alpha(\mu_\alpha \otimes I_\alpha)(h \otimes k \otimes b) \\
&= \omega_\alpha(\mu_\alpha \otimes I_\alpha)(h \otimes k \otimes b)
\end{aligned}$$

and

$$\omega_\alpha(\eta_\alpha \otimes I_\alpha)(k \otimes b) = \omega_\alpha(\eta_\alpha(k) \otimes b) = \sigma_\alpha(\eta_\alpha(k))b = kb =\sim (k \otimes b)$$

Now, we'll prove that σ_α is left H_α-module map $\forall \alpha \in \pi$ i.e., the following diagram is commute

$$\begin{tikzcd}
H_\alpha \arrow[r, "\sigma_\alpha"] & C_\alpha \\
H_\alpha \otimes H_\alpha \arrow[u, "\mu_\alpha^H"] \arrow[r, "I_\alpha \otimes \sigma_\alpha"'] & H_\alpha \otimes C_\alpha \arrow[u, "\omega_\alpha"']
\end{tikzcd}$$

$$\omega_\alpha(I_\alpha \otimes \sigma_\alpha) = \mu_\alpha^C(\sigma_\alpha \otimes I_\alpha)(I_\alpha \otimes \sigma_\alpha) = \mu_\alpha^C(\sigma_\alpha \otimes \sigma_\alpha) = \sigma_\alpha\mu_\alpha^H.$$

∎

Remark 2.2. If (C, σ) is a Hopf $\pi-$subcoalgebra of H, then the map $L_{\alpha,\beta} = (\sigma_\alpha \otimes I^H_\beta)\Delta^H_{\alpha,\beta} : H_{\alpha\beta} \to C_\alpha \otimes H_\beta \ \ \forall \alpha, \beta \in \pi$ is an algebra map.

Proposition 2.3. *Let $H = \{H_\alpha\}_{\alpha\in\pi}$ be a Hopf $\pi-$coalgebra and (C, σ) be a Hopf $\pi-$subcoalgebra of H. Let $B = \{B_\alpha\}_{\alpha\in\pi}$, where $B_\alpha = \{h \in H_\alpha : L_{1,\alpha}(h) = 1 \otimes h\}$, then B_α is subalgebra of H_α and B is right $\pi-$coideal of H (this B is called $\pi-$quantum right embeddable homogeneous space of H).*

Proof. Firstly, let $h, k \in B_\alpha$, then $L_{1,\alpha}(hk) = L_{1,\alpha}(h) \cdot L_{1,\alpha}(k) = (1 \otimes h) \cdot (1 \otimes k) = 1 \otimes hk$ and hence $hk \in B_\alpha$. Secondly let $h \in B_{\alpha\beta}$, we will prove that $\Delta^H_{\alpha,\beta}(h) \in B_\alpha \otimes H_\beta$.

$$
\begin{aligned}
&(L_{1,\alpha} \otimes I^H_\beta)\Delta^H_{\alpha,\beta}(h)\\
&= ((\sigma_1 \otimes I^H_\alpha)\Delta^H_{1,\alpha} \otimes I^H_\beta)\Delta^H_{\alpha,\beta}(h)\\
&= (\sigma_1 \otimes I^H_\alpha \otimes I^H_\beta)(\Delta^H_{1,\alpha} \otimes I^H_\beta)\Delta^H_{\alpha,\beta}(h)\\
&= (\sigma_1 \otimes I^H_\alpha \otimes I^H_\beta)(I^H_1 \otimes \Delta^H_{\alpha,\beta})\Delta^H_{1,\alpha\beta}(h)\\
&= (I^C_1 \otimes \Delta^H_{\alpha,\beta})(\sigma_1 \otimes I^H_{\alpha\beta})\Delta^H_{1,\alpha\beta}(h)\\
&= (I^C_1 \otimes \Delta^H_{\alpha,\beta})(1 \otimes h)\\
&= 1 \otimes \Delta^H_{\alpha,\beta}(h).
\end{aligned}
$$

∎

Lemma 2.4. *Let $H = \{H_\alpha\}_{\alpha\in\pi}$ be a Hopf $\pi-$coalgebra and (C, σ) be a left $\pi-$coisotropic quantum subgroup of H. For $\alpha, \beta, \gamma \in \pi$, $a, b \in H_{\alpha\beta}$ we have*

- $L_{\alpha,\beta}(ab) = \Delta^H_{\alpha,\beta}(a)\Theta L_{\alpha,\beta}(b)$
- $(I_\alpha \otimes \Delta^H_{\beta,\gamma})L_{\alpha,\beta\gamma} = (L_{\alpha,\beta} \otimes I_\gamma)\Delta^H_{\alpha\beta,\gamma}$,

where $(m \otimes n)\Theta(u \otimes v) = \omega_\alpha(m \otimes u) \otimes nv, m \in H_\alpha, u \in C_\alpha, n, v \in H_\beta$.

Proof.

$$
\begin{aligned}
&L_{\alpha,\beta}(ab)\\
&= (\sigma_\alpha \otimes I^H_\beta)\Delta^H_{\alpha,\beta}\mu_{\alpha\beta}(a \otimes b)\\
&= (\sigma_\alpha \otimes I^H_\beta)(\mu_\alpha \otimes \mu_\beta)(I \otimes \tau \otimes I)(\Delta^H_{\alpha,\beta} \otimes \Delta^H_{\alpha,\beta})(a \otimes b)\\
&= (\sigma_\alpha\mu_\alpha \otimes \mu_\beta)(I \otimes \tau \otimes I)(\Delta^H_{\alpha,\beta} \otimes \Delta^H_{\alpha,\beta})(a \otimes b)\\
&= \omega_\alpha(a^\alpha_1 \otimes \sigma_\alpha(b^\alpha_1)) \otimes a^\beta_2 b^\beta_2\\
&= \Delta^H_{\alpha,\beta}(a)\Theta L_{\alpha,\beta}(b)
\end{aligned}
$$

Also,

$$\begin{aligned}
&(I_\alpha \otimes \Delta^H_{\beta,\gamma})L_{\alpha,\beta\gamma}\\
=\ &(I_\alpha \otimes \Delta^H_{\beta,\gamma})(\sigma_\alpha \otimes I^H_{\beta\gamma})\Delta^H_{\alpha,\beta\gamma}\\
=\ &(\sigma_\alpha \otimes I^H_\beta \otimes I^H_\gamma)(I_\alpha \otimes \Delta^H_{\beta,\gamma})\Delta^H_{\alpha,\beta\gamma}\\
=\ &(\sigma_\alpha \otimes I^H_\beta \otimes I^H_\gamma)(\Delta^H_{\alpha,\beta} \otimes I_\gamma)\Delta^H_{\alpha\beta,\gamma}\\
=\ &((\sigma_\alpha \otimes I^H_\beta)\Delta^H_{\alpha,\beta} \otimes I^H_\gamma)\Delta^H_{\alpha\beta,\gamma}\\
=\ &(L_{\alpha,\beta} \otimes I_\gamma)\Delta^H_{\alpha\beta,\gamma}
\end{aligned}$$

∎

Proposition 2.5. *Let $H = \{H_\alpha\}_{\alpha\in\pi}$ be a Hopf $\pi-$coalgebra and let (C, σ) be a left $\pi-$coisotropic quantum subgroup of H. Let $G = \{G_\alpha\}_{\alpha\in\pi}$, where $G_\alpha = \{h \in H_\alpha : L_{1,\alpha}(h) = \sigma_1(1) \otimes h\}$, then G_α is subalgebra of H_α and G is right $\pi-$ coideal of H.*

Proof. For $h, g \in G_\alpha$, we have $L_{1,\alpha}(h) = \sigma_1(1) \otimes h$, $L_{1,\alpha}(g) = \sigma_1(1) \otimes g$ and $\omega_1(I_1 \otimes \sigma_1) = \sigma_1\mu_1$.
We‘ll prove that $hg \in G_\alpha, i.e., L_{1,\alpha}(hg) = \sigma_1(1) \otimes hg$

$$\begin{aligned}
&L_{1,\alpha}(hg)\\
=\ &\Delta_{1,\alpha}(h)\Theta L_{1,\alpha}(g)\\
=\ &\Delta_{1,\alpha}(h)\Theta(\sigma_1(1) \otimes g)\\
=\ &(\omega_1 \otimes \mu_\alpha)(h^1_1 \otimes \sigma_1(1) \otimes h^\alpha_2 \otimes g)\\
=\ &\omega_1(I_1 \otimes \sigma_1)(h^1_1 \otimes 1) \otimes h^\alpha_2 g\\
=\ &\sigma_1\mu_1(h^1_1 \otimes 1) \otimes h^\alpha_2 g\\
=\ &\sigma_1(h^1_1) \otimes h^\alpha_2 g\\
=\ &(\sigma_1 \otimes \mu_\alpha)(\Delta^H_{1,\alpha} \otimes I_\alpha)(h \otimes g)\\
=\ &(I_1 \otimes \mu_\alpha)((\sigma_1 \otimes I_1)\Delta^H_{1,\alpha}(h) \otimes g)\\
=\ &(I_1 \otimes \mu_\alpha)(\sigma_1(1) \otimes h \otimes g)\\
=\ &\sigma_1(1) \otimes hg
\end{aligned}$$

We‘ll prove that G is right π-coideal of H, i.e., for $h \in G_{\alpha\beta}$, $\Delta^H_{\alpha,\beta}(h) \in G_\alpha \otimes H_\beta$

$$\begin{aligned}
&(L_{1,\alpha} \otimes I^H_\beta)\Delta^H_{\alpha,\beta}(h) \\
&= ((\sigma_1 \otimes I^H_\alpha)\Delta^H_{1,\alpha} \otimes I^H_\beta)\Delta^H_{\alpha,\beta}(h) \\
&= (\sigma_1 \otimes I^H_\alpha \otimes I^H_\beta)(\Delta^H_{1,\alpha} \otimes I^H_\beta)\Delta^H_{\alpha,\beta}(h) \\
&= (\sigma_1 \otimes I^H_\alpha \otimes I^H_\beta)(I^H_1 \otimes \Delta^H_{\alpha,\beta})\Delta^H_{1,\alpha\beta}(h) \\
&= (I^C_1 \otimes \Delta^H_{\alpha,\beta})(\sigma_1 \otimes I^H_{\alpha\beta})\Delta^H_{1,\alpha\beta}(h) \\
&= (I^C_1 \otimes \Delta^H_{\alpha,\beta})(\sigma_1(1) \otimes h) \\
&= \sigma_1(1) \otimes \Delta^H_{1,\alpha\beta}(h)
\end{aligned}$$

∎

Theorem 2.6. *Suppose* $V = \{V_\alpha\}_{\alpha\in\pi}$, *be a right* $\pi-$*comodule over the left* $\pi-$*coisotropic quantum subgroup* C *of Hopf* $\pi-$*coalgebra* H *by* $\rho = \{\rho_{\alpha,\beta} : V_{\alpha\beta} \to V_\alpha \otimes C_\beta\}_{\alpha,\beta\in\pi}$. *Then* $Ind(\rho) = \{Ind(\rho)_\alpha\}_{\alpha\in\pi}$ *where* $Ind(\rho)_\alpha = \{x \in V_1 \otimes H_\alpha : (I_1 \otimes L_{1,\alpha})x = (\rho_{1,1} \otimes I_\alpha)x\}$ *is right* $\pi-$*comodule over* H *by* $(I\otimes\Delta) = \{(I \otimes \Delta)_{\alpha,\beta} = I_1 \otimes \Delta_{\alpha,\beta}\}_{\alpha,\beta\in\pi}$

Proof. We will prove that, for $\alpha, \beta \in \pi$, $(I\otimes\Delta)_{\alpha,\beta}(Ind(\rho)_{\alpha\beta}) \subseteq Ind(\rho)_\alpha \otimes H_\beta$. Since, for $v \otimes h \in Ind(\rho)_{\alpha\beta}$, we have

$$\begin{aligned}
&(I_1 \otimes L_{1,\alpha} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})(v \otimes h) \\
&= (I_1 \otimes (L_{1,\alpha} \otimes I_\beta)\Delta_{\alpha,\beta})(v \otimes h) \\
&= (I_1 \otimes (I_1 \otimes \Delta_{\alpha,\beta})L_{1,\alpha\beta})(V \otimes h) && \text{by Lemma 2.4} \\
&= (I_1 \otimes I_1 \otimes \Delta_{\alpha,\beta})(I_1 \otimes L_{1,\alpha\beta})(v \otimes h) \\
&= (I_1 \otimes I_1 \otimes \Delta_{\alpha,\beta})(\rho_{1,1} \otimes I_{\alpha\beta})(v \otimes h) \\
&= (\rho_{1,1} \otimes I_\alpha \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})(v \otimes h)
\end{aligned}$$

Now, we‘ll prove that the following diagrams are commute

$$\begin{tikzcd}
Ind(\rho)_{\alpha\beta\gamma} \arrow[r, "(I\otimes\Delta)_{\alpha\beta,\gamma}"] \arrow[d, "(I\otimes\Delta)_{\alpha,\beta\gamma}"'] & Ind(\rho)_{\alpha\beta} \otimes H_\gamma \arrow[d, "(I\otimes\Delta)_{\alpha,\beta} \otimes I_\gamma"] \\
Ind(\rho)_\alpha \otimes H_{\beta\gamma} \arrow[r, "I_{Ind(\rho)_\alpha} \otimes \Delta_{\beta,\gamma}"'] & Ind(\rho)_\alpha \otimes H_\beta \otimes H_\gamma
\end{tikzcd}$$

and

$$\begin{tikzcd}
Ind(\rho)_\alpha \arrow[rr, "(I\otimes\Delta)_{\alpha,1}"] \arrow[dr, "\sim_{Ind(\rho)_\alpha}"'] & & Ind(\rho)_\alpha \otimes H_1 \arrow[dl, "I_{Ind(\rho)_\alpha} \otimes \epsilon^H"] \\
& Ind(\rho)_\alpha \otimes K &
\end{tikzcd}$$

$$
\begin{aligned}
& ((I \otimes \Delta)_{\alpha,\beta} \otimes I_\gamma)(I \otimes \Delta)_{\alpha\beta,\gamma} \\
&= (I_1 \otimes \Delta_{\alpha,\beta} \otimes I_\gamma)(I_1 \otimes \Delta_{\alpha\beta,\gamma}) \\
&= (I_1 \otimes (\Delta_{\alpha,\beta} \otimes I_\gamma)\Delta_{\alpha\beta,\gamma}) \\
&= (I_1 \otimes (I_\alpha \otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma}) \\
&= (I_1 \otimes I_\alpha \otimes \Delta_{\beta,\gamma})(I_1 \otimes \Delta_{\alpha,\beta\gamma}) \\
&= (I_{Ind(\rho)_\alpha} \otimes \Delta_{\beta,\gamma})(I \otimes \Delta)_{\alpha,\beta\gamma}
\end{aligned}
$$

and

$$
\begin{aligned}
& (I_{Ind(\rho)_\alpha} \otimes \epsilon^H)(I \otimes \Delta)_{\alpha,1} \\
&= (I_1 \otimes I_\alpha \otimes \epsilon^H)(I_1 \otimes \Delta_{\alpha,1}) \\
&= (I_1 \otimes (I_\alpha \otimes \epsilon)\Delta_{\alpha,1}) \\
&= (I_1 \otimes \sim_{H_\alpha}) \\
&= \sim_{Ind(\rho)_\alpha}
\end{aligned}
$$

$\blacksquare$

Remark 2.7. Given a right corepresentation ρ of the $\pi-$coisotropic quantum subgroup of (C, σ) the corresponding corepresentation $(I \otimes \Delta)$ on $Ind(\rho)$ of H is called induced representation from ρ on H.

3 Geometric realization for induced representation

In this section we given the geometric interperation of the induced representation. Throughout this section H is Hopf $\pi-$ coalgebra, (C, σ) is Hopf $\pi-$subcoalgebra of H and $V = \{V_\alpha\}_{\alpha \in \pi}$ be a right $\pi-$comodule over C. The purpose of this section is to show that the induced representation $Ind(\rho)$ from Hopf group subcoalgebra H is isomorphic as module to the tensor product of $\pi-$quantum embeddable homogeneous space B (in Proposition 2.3) and the given comodule V. In case (C, σ) is left $\pi-$coisotropic quantum subgroup, then H is isomorphic to $C \otimes G$ as a vector space where $G = \{G_\alpha\}_{\alpha \in \pi}$, where

$$G_\alpha = \{h \in H_\alpha : L_{1,\alpha}(h) = (\sigma_1 \otimes I_\alpha^H)\Delta_{1,\alpha}^H(h) = \sigma_1(1) \otimes h\}.$$

Lemma 3.1. *$Ind(\rho)_\alpha$ is a right (left) B_α-module for all $\alpha \in \pi$.*

Proof. Let $v_1 \otimes h_\alpha \in Ind(\rho)_\alpha, b_\alpha \in B_\alpha$, we define the right action as follows $\lambda_\alpha(v_1 \otimes h_\alpha \otimes b_\alpha) = v_1 \otimes h_\alpha b_\alpha$. We need only to prove that $\lambda_\alpha(v_1 \otimes h_\alpha \otimes b_\alpha) = v_1 \otimes b_\alpha h_\alpha \in Ind(\rho)_\alpha$. We have $(I_1 \otimes L_{1,\alpha})(v_1 \otimes h_\alpha) = (\rho_{1,1} \otimes I_\alpha)(v_1 \otimes h_\alpha)$

imply that$((I_1\otimes L_{1,\alpha})(v_1\otimes h_\alpha))(1\otimes 1\otimes b_\alpha) = ((\rho_{1,1}\otimes I_\alpha)(v_1\otimes h_\alpha))(1\otimes 1\otimes b_\alpha)$ imply that

$$v_1\otimes\sigma_1((h_\alpha)_1)\otimes(h_\alpha)_2b_\alpha = \rho_{1,1}(v_1)\otimes h_\alpha b_\alpha. \qquad (*)$$

Also, we have

$$\begin{aligned}
&L_{1,\alpha}(h_\alpha b_\alpha)\\
&= \Delta^H_{1,\alpha}(h_\alpha)\Theta L_{1,\alpha}(b_\alpha)\\
&= ((h_\alpha)_1\otimes(h_\alpha)_2)\Theta(1_{C_1}\otimes b_\alpha) && \text{by lemma 2.5}\\
&= \omega_\alpha((h_\alpha)_1\otimes\sigma_1(1_{H_1}))\otimes(h_\alpha)_2b_\alpha\\
&= \sigma_1(h_{\alpha_1})\otimes(h_\alpha)_2b_\alpha. && (**)
\end{aligned}$$

Now,

$$\begin{aligned}
&(I_1\otimes L_{1,\alpha})\lambda_\alpha(v_1\otimes h_\alpha\otimes b_\alpha)\\
&= (I_1\otimes L_{1,\alpha})(v_1\otimes h_\alpha b_\alpha)\\
&= v_1\otimes L_{1,\alpha}(h_\alpha b_\alpha)\\
&= v_1\otimes\sigma_1((h_\alpha)_1)\otimes(h_\alpha)_2b_\alpha && \text{by } (**)\\
&= \rho_{1,1}(v_1)\otimes h_\alpha b_\alpha && \text{by } (*)\\
&= (\rho_{1,1}\otimes I_\alpha)(v_1\otimes h_\alpha b_\alpha)\\
&= (\rho_{1,1}\otimes I_\alpha)\lambda_\alpha(v_1\otimes h_\alpha\otimes b_\alpha).
\end{aligned}$$

The left action is similar. ∎

Definition 3.2. Let (C,σ) be a Hopf $\pi-$subcoalgebra of H. If $g=\{g_\alpha : C_\alpha\to H_\alpha\}_{\alpha\in\pi}$, is a family of linear maps, its convolution inverse, if it exists, is a family of linear maps $g^{-1}=\{g_\alpha^{-1}: C_{\alpha^{-1}}\to H_\alpha\}_{\alpha\in\pi}$ such that

$$\mu_\alpha(g_\alpha\otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}(c) = \epsilon^C(c)1_{H_\alpha} = \mu_\alpha(g_\alpha^{-1}\otimes g_\alpha)\Delta^C_{\alpha^{-1},\alpha}(c)$$

Definition 3.3. A Hopf $\pi-$subcoalgebra (C,σ) of H is said to have a left section if there exists a family of linear, convolution invertible, maps $g=\{g_\alpha : C_\alpha\to H_\alpha\}_{\alpha\in\pi}$ such that for all $\alpha\in\pi$, $g_\alpha(1)=1$ and $L_{1,\alpha}g_\alpha=(I_1\otimes g_\alpha)\Delta^C_{1,\alpha}$

Lemma 3.4. *For any Hopf $\pi-$coalgebra H we have*

1. $(\Delta^H_{1,\alpha}\otimes\Delta^H_{\alpha^{-1},1})\Delta^H_{\alpha,\alpha^{-1}} = (I_1\otimes\Delta^H_{\alpha,\alpha^{-1}}\otimes I_1)(\Delta^H_{1,1}\otimes I_1)\Delta^H_{1,1}$
2. $(\Delta^H_{\alpha^{-1},1}\otimes\Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha} = (I_{\alpha^{-1}}\otimes\Delta^H_{1,1}\otimes I_\alpha)(\Delta^H_{\alpha^{-1},1}\otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}$

Proof.

$$\begin{aligned}
&(\Delta^H_{1,\alpha}\otimes\Delta^H_{\alpha^{-1},1})\Delta^H_{\alpha,\alpha^{-1}}\\
&=(\Delta^H_{1,\alpha}\otimes I_{\alpha^{-1}}\otimes I_1)(I_\alpha\otimes\Delta^H_{\alpha^{-1},1})\Delta^H_{\alpha,\alpha^{-1}}\\
&=(\Delta^H_{1,\alpha}\otimes I_{\alpha^{-1}}\otimes I_1)(\Delta^H_{\alpha,\alpha^{-1}}\otimes I_1)\Delta^H_{1,1}\\
&=((\Delta^H_{1,\alpha}\otimes I_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}}\otimes I_1)\Delta^H_{1,1}\\
&=((I_1\otimes\Delta^H_{\alpha,\alpha^{-1}})\Delta^H_{1,1}\otimes I_1)\Delta^H_{1,1}\\
&=(I_1\otimes\Delta^H_{\alpha,\alpha^{-1}}\otimes I_1)(\Delta^H_{1,1}\otimes I_1)\Delta^H_{1,1}
\end{aligned}$$

$$i.e., (\Delta^H_{1,\alpha}\otimes\Delta^H_{\alpha^{-1},1})\Delta^H_{\alpha,\alpha^{-1}}(h)=h^1_{11}\otimes h^\alpha_{121}\otimes h^{\alpha^{-1}}_{122}\otimes h^1_2$$

The second statement is similar. ∎

Now, from lemma 3.5 up to theorem 3.12, H is Hopf $\pi-$coalgebra and (C,σ) is Hopf $\pi-$subcoalgebra. Also, (C,σ) have a section $g=\{g_\alpha\}_{\alpha\in\pi}$ and antipode $S^C=\{S^C_\alpha : C_\alpha\to C_{\alpha^{-1}}\}_{\alpha\in\pi}$

Lemma 3.5. *For* $\alpha\in\pi, L_{1,\alpha}g_\alpha^{-1}=(S^C_1\otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}$

Proof. We will prove that for $\alpha\in\pi$, $L_{1,\alpha}g_\alpha^{-1}$ and $(S^C_1\otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}$ are inverse to the same element $L_{1,\alpha}g_\alpha$ in the convolution algebra $Conv(C,C_1\otimes H_\alpha)$. For $c\in C_1$,

$$\begin{aligned}
&(L_{1,\alpha}g_\alpha * L_{1,\alpha}g_\alpha^{-1})(c)\\
&=\{(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha * (\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha^{-1}\}(c)\\
&=\mu_{C_1\otimes H_\alpha}\{(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha\otimes(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha^{-1}\}\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&=(\mu_{C_1}\otimes\mu_{H_\alpha})(I\otimes\tau\otimes I)\{(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha\otimes(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha^{-1}\}\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&=(\mu_{C_1}\otimes\mu_{H_\alpha})(\sigma_1\otimes\sigma_1\otimes I^H_\alpha\otimes I^H_\alpha)(I\otimes\tau\otimes I)(\Delta^H_{1,\alpha}\otimes\Delta^H_{1,\alpha})(g_\alpha\otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&=(\mu_{C_1}(\sigma_1\otimes\sigma_1)\otimes\mu_{H_\alpha})(I\otimes\tau\otimes I)(\Delta^H_{1,\alpha}\otimes\Delta^H_{1,\alpha})(g_\alpha\otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&=(\sigma_1\mu_{H_1}\otimes\mu_{H_\alpha})(I\otimes\tau\otimes I)(\Delta^H_{1,\alpha}\otimes\Delta^H_{1,\alpha})(g_\alpha\otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&=(\sigma_1\otimes I^H_\alpha)(\mu_{H_1}\otimes\mu_{H_\alpha})(I\otimes\tau\otimes I)(\Delta^H_{1,\alpha}\otimes\Delta^H_{1,\alpha})(g_\alpha\otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&=(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}\mu_{H_\alpha}(g_\alpha\otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&=(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}(\epsilon^C(c)1_{H_\alpha})\\
&=\epsilon^C(c)(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}(1_{H_\alpha})\\
&=\epsilon^C(c)(1_{C_1}\otimes 1_{H_\alpha}).
\end{aligned}$$

and

$$
\begin{aligned}
&\{L_{1,\alpha}g_\alpha * (S_1^C \otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}\}(c)\\
&= \{(\sigma_1 \otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha * (S_1^C \otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}\}(c)\\
&= \{(I_1 \otimes g_\alpha)\Delta^C_{1,\alpha} * (S_1^C \otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}\}(c)\\
&= \mu_{C_1\otimes H_\alpha}\{(I_1 \otimes g_\alpha)\Delta^C_{1,\alpha} \otimes (S_1^C \otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}\}\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&= (\mu_{C_1} \otimes \mu_{H_\alpha})(I \otimes \tau \otimes I)\{(I_1 \otimes g_\alpha)\Delta^C_{1,\alpha} \otimes (S_1^C \otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}\}\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&= (\mu_{C_1} \otimes \mu_{H_\alpha})(I \otimes \tau \otimes I)(I_1 \otimes g_\alpha \otimes S_1^C \otimes g_\alpha^{-1})(I_1 \otimes I_\alpha \otimes \tau)(\Delta^C_{1,\alpha} \otimes \Delta^C_{\alpha^{-1},1})\Delta^C_{\alpha,\alpha^{-1}}(c)\\
&= (\mu_{C_1} \otimes \mu_{H_\alpha})(I \otimes \tau \otimes I)(I_1 \otimes g_\alpha \otimes S_1^C \otimes g_\alpha^{-1})(I_1 \otimes I_\alpha \otimes \tau)(c^1_{11} \otimes c^\alpha_{121} \otimes c^{\alpha^{-1}}_{122} \otimes c^1_2)\\
&= c^1_{11}S_1^C(c^1_2) \otimes g_\alpha(c^\alpha_{121})g_\alpha^{-1}(c^{\alpha^{-1}}_{122})\\
&= c^1_{11}S_1^C(c^1_2) \otimes \epsilon^C(c^\alpha_{12})1_{H_\alpha}\\
&= \epsilon^C(c^\alpha_{12})c^1_{11}S_1^C(c^1_2) \otimes 1_{H_\alpha}\\
&= c^1_1S_1^C(c^1_2) \otimes 1_{H_\alpha}\\
&= \epsilon^C(c)(1_{C_1} \otimes 1_{H_\alpha})
\end{aligned}
$$

∎

Lemma 3.6. *For $h \in H_\alpha$, we have $h = g_\alpha\sigma_\alpha(h^\alpha_1)g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{21})h^\alpha_{22}$.*

Proof.

$$
\begin{aligned}
h &= \epsilon^H(h^1_1)h^\alpha_2\\
&= \varepsilon^C\sigma_1(h^1_1)h^\alpha_2\\
&= (\mu_\alpha(g_\alpha \otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}(\sigma_1(h^1_1)))h^\alpha_2\\
&= (\mu_\alpha(g_\alpha \otimes g_\alpha^{-1})(\sigma_\alpha \otimes \sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}}(h^1_1))h^\alpha_2\\
&= \mu_\alpha(\mu_\alpha(g_\alpha \otimes g_\alpha^{-1})(\sigma_\alpha \otimes \sigma_{\alpha^{-1}}) \otimes I_\alpha)(\Delta^H_{\alpha,\alpha^{-1}} \otimes I_\alpha)\Delta^H_{1,\alpha}(h)\\
&= \mu_\alpha(\mu_\alpha(g_\alpha \otimes g_\alpha^{-1})(\sigma_\alpha \otimes \sigma_{\alpha^{-1}}) \otimes I_\alpha)(I_\alpha \otimes \Delta^H_{\alpha^{-1},\alpha})\Delta^H_{\alpha,1}(h)\\
&= g_\alpha\sigma_\alpha(h^\alpha_1)g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{21})h^\alpha_{22}.
\end{aligned}
$$

∎

Lemma 3.7. *For $h \in H_1$ and $\alpha \in \pi$, we have $\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha)\Delta^H_{\alpha,\alpha^{-1}}(h) \in B_\alpha$.*

Proof.

$$\begin{aligned}
&L_{1,\alpha}\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}}\otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h)\\
&=(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}}\otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h)\\
&=\mu_{C_1\otimes H_\alpha}((\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha^{-1}\sigma_{\alpha^{-1}}\otimes(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h)\\
&=\mu_{C_1\otimes H_\alpha}((S^C_1\otimes g_\alpha^{-1})\tau\Delta^C_{\alpha^{-1},1}\sigma_{\alpha^{-1}}\otimes(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h)\\
&=\mu_{C_1\otimes H_\alpha}((S^C_1\otimes g_\alpha^{-1})\tau(\sigma_{\alpha^{-1}}\otimes\sigma_1)\Delta^H_{\alpha^{-1},1}\otimes(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h)\\
&=\mu_{C_1\otimes H_\alpha}((S^C_1\otimes g_\alpha^{-1})\tau\otimes I_1\otimes I^H_\alpha)(\sigma_{\alpha^{-1}}\otimes\sigma_1\otimes\sigma_1\otimes I^H_\alpha)(\Delta^H_{\alpha^{-1},1}\otimes\Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h)\\
&=\mu_{C_1\otimes H_\alpha}((S^C_1\otimes g_\alpha^{-1})\tau\otimes I_1\otimes I^H_\alpha)(\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{11})\otimes\sigma_1(h^1_{121})\otimes\sigma_1(h^1_{122})\otimes h^\alpha_2)\\
&=S^C_1(\sigma_1(h^1_{121}))\sigma_1(h^1_{122})\otimes g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{11})h^\alpha_2\\
&=\epsilon^C(\sigma_1(h^1_{12})1\otimes g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{11})h^\alpha_2\\
&=\epsilon^H(h^1_{12})1\otimes g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{11})h^\alpha_2\\
&=1\otimes g_\alpha^{-1}\sigma_{\alpha^{-1}}(\epsilon^H(h^1_{12})h^{\alpha^{-1}}_{11})h^\alpha_2\\
&=1\otimes g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{1})h^\alpha_2\\
&=1\otimes\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}}\otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h).
\end{aligned}$$

∎

Lemma 3.8. *For $\alpha\in\pi$,*

$$(\sigma_\alpha\otimes\sigma_{\alpha^{-1}}\otimes I^H_\alpha)(\Delta^H_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha=(I_\alpha\otimes I_{\alpha^{-1}}\otimes g_\alpha)(I_\alpha\otimes\Delta^C_{\alpha^{-1},\alpha})\Delta^C_{\alpha,1}$$

Proof.

$$\begin{aligned}
&(\sigma_\alpha\otimes\sigma_{\alpha^{-1}}\otimes I^H_\alpha)(\Delta^H_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha\\
&=((\sigma_\alpha\otimes\sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha\\
&=(\Delta^C_{\alpha,\alpha^{-1}}\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha\\
&=(\Delta^C_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}g_\alpha\\
&=(\Delta^C_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)(I_1\otimes g_\alpha)\Delta^C_{1,\alpha}\\
&=(I_\alpha\otimes I_{\alpha^{-1}}\otimes g_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes I_\alpha)\Delta^C_{1,\alpha}\\
&=(I_\alpha\otimes I_{\alpha^{-1}}\otimes g_\alpha)(I_\alpha\otimes\Delta^C_{\alpha^{-1},\alpha})\Delta^C_{\alpha,1}
\end{aligned}$$

∎

Lemma 3.9. *For $\alpha\in\pi$ and $b\in B_\alpha$*

$$(\sigma_\alpha\otimes\sigma_{\alpha^{-1}}\otimes I^H_\alpha)(\Delta^H_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)\Delta^H_{1,\alpha}(b)=1\otimes1\otimes b$$

Proof.

$$\begin{aligned}
&(\sigma_\alpha \otimes \sigma_{\alpha^{-1}} \otimes I^H_\alpha)(\Delta^H_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{1,\alpha}(b)\\
&= ((\sigma_\alpha \otimes \sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{1,\alpha}(b)\\
&= (\Delta^C_{\alpha,\alpha^{-1}}\sigma_1 \otimes I^H_\alpha)\Delta^H_{1,\alpha}(b)\\
&= (\Delta^C_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)(\sigma_1 \otimes I^H_\alpha)\Delta^H_{1,\alpha}(b)\\
&= (\Delta^C_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)(1 \otimes b)\\
&= 1 \otimes 1 \otimes b
\end{aligned}$$

∎

Theorem 3.10. *H is isomorphic to $C \otimes B$ as vector space.*

Proof. We define $A = \{A_\alpha : C_\alpha \otimes B_\alpha \to H_\alpha\}_{\alpha\in\pi}$ as follow

$$A_\alpha(c \otimes b) = \mu_\alpha(g_\alpha \otimes I_\alpha)(c \otimes b) = g_\alpha(c)b.$$

Clear A_α is linear and by lemma 3.6 and lemma 3.7 that A_α is surjective for all $\alpha \in \pi$. We define $A^{-1}_\alpha = (\sigma_\alpha \otimes \mu_\alpha(g^{-1}_\alpha\sigma_{\alpha^{-1}} \otimes I_\alpha))(\Delta^H_{\alpha,\alpha^{-1}} \otimes I_\alpha)\Delta^H_{1,\alpha}$. We'll prove that for $\alpha \in \pi, A_\alpha A^{-1}_\alpha = I_{H_\alpha}$ and $A^{-1}_\alpha A_\alpha = I_{C_\alpha \otimes B_\alpha}$. Firstly, let $h \in H_\alpha$ and $c \otimes b \in C_\alpha \otimes B_\alpha$

$$\begin{aligned}
&A_\alpha A^{-1}_\alpha(h)\\
&= A_\alpha((\sigma_\alpha \otimes \mu_\alpha(g^{-1}_\alpha\sigma_{\alpha^{-1}} \otimes I_\alpha))(\Delta^H_{\alpha,\alpha^{-1}} \otimes I_\alpha)\Delta^H_{1,\alpha}(h))\\
&= A_\alpha((\sigma_\alpha \otimes \mu_\alpha(g^{-1}_\alpha\sigma_{\alpha^{-1}} \otimes I_\alpha))(I_\alpha \otimes \Delta^H_{\alpha^{-1},\alpha})\Delta^H_{\alpha,1}(h))\\
&= A_\alpha(\sigma_\alpha(h^\alpha_1) \otimes g^{-1}_\alpha\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{21})h^\alpha_{22})\\
&= g_\alpha\sigma_\alpha(h^\alpha_1)g^{-1}_\alpha\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{21})h^\alpha_{22}\\
&= h \qquad\qquad \text{by lemma 3.6}
\end{aligned}$$

Secondly, since, for $\alpha \in \pi$ we have $((\sigma_\alpha \otimes \sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{1,\alpha}$ is an algebra map, then

$$\begin{aligned}
& A_\alpha^{-1} A_\alpha(c \otimes b) \\
&= (\sigma_\alpha \otimes \mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha))(\Delta^H_{\alpha,\alpha^{-1}} \otimes I_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha \otimes I_\alpha)(c \otimes b) \\
&= (I_\alpha \otimes \mu_\alpha)(I_\alpha \otimes g_\alpha^{-1} \otimes I_\alpha)(\sigma_\alpha \otimes \sigma_{\alpha^{-1}} \otimes I^H_\alpha)(\Delta^H_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha \otimes I_\alpha)(c \otimes b) \\
&= (I_\alpha \otimes \mu_\alpha)(I_\alpha \otimes g_\alpha^{-1} \otimes I_\alpha)((\sigma_\alpha \otimes \sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha \otimes I_\alpha)(c \otimes b) \\
&= (I_\alpha \otimes \mu_\alpha)(I_\alpha \otimes g_\alpha^{-1} \otimes I_\alpha)\mu_{H_\alpha \otimes H_{\alpha^{-1}} \otimes H_\alpha} \\
&\quad (((\sigma_\alpha \otimes \sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{1,\alpha} g_\alpha \otimes ((\sigma_\alpha \otimes \sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{1,\alpha})(c \otimes b) \\
&= (I_\alpha \otimes \mu_\alpha)(I_\alpha \otimes g_\alpha^{-1} \otimes I_\alpha)\mu_{H_\alpha \otimes H_{\alpha^{-1}}} \\
&\quad ((I_\alpha \otimes I_{\alpha^{-1}} \otimes g_\alpha)(I_\alpha \otimes \Delta^C_{\alpha^{-1},\alpha})\Delta^C_{\alpha,1}(c) \otimes 1 \otimes 1 \otimes b) \qquad \text{by lemmas 3.8, 3.9} \\
&= c_1^\alpha \otimes g_\alpha^{-1}(c_{21}^{\alpha^{-1}}) g_\alpha(c_{22}^\alpha) b \\
&= c_1^\alpha \otimes \epsilon(c_2^1) b \\
&= \epsilon(c_2^1) c_1^\alpha \otimes b \\
&= c \otimes b.
\end{aligned}$$

∎

Remark 3.11. In any Hopf π -coalgebra H, every H_α is left H_1 -comodule by $\Delta_{1,\alpha}$. i.e., the following diagrams are commute

$$\begin{tikzcd}
H_\alpha \arrow[r, "\Delta_{1,\alpha}"] \arrow[d, "\Delta_{1,\alpha}"'] & H_1 \otimes H_\alpha \arrow[d, "\Delta_{1,1} \otimes I_\alpha"] \\
H_1 \otimes H_\alpha \arrow[r, "I_1 \otimes \Delta_{1,\alpha}"'] & H_1 \otimes H_1 \otimes H_\alpha
\end{tikzcd}
\qquad
\begin{tikzcd}
H_\alpha \arrow[r, "\Delta_{1,\alpha}"] \arrow[dr, "\sim"'] & H_1 \otimes H_\alpha \arrow[d, "\epsilon^H \otimes I_\alpha"] \\
 & K \otimes H_\alpha
\end{tikzcd}$$

A Hopf $\pi-$coalgebra H is called have a left (right) cosection if there exist a family of algebra maps $\eta = \{\eta_\alpha : H_1 \to H_\alpha\}_{\alpha \in \pi}$ such that η_α is left (right) H_1-comodule map $\forall \alpha \in \pi$ i.e., the following diagram is commute

$$\begin{tikzcd}
H_1 \arrow[r, "\eta_\alpha"] \arrow[d, "\Delta_{1,1}"'] & H_\alpha \arrow[d, "\Delta_{1,\alpha}"] \\
H_1 \otimes H_1 \arrow[r, "I_1 \otimes \eta_\alpha"'] & H_1 \otimes H_\alpha
\end{tikzcd}$$

Theorem 3.12. *If H have a left cosection, then $Ind(\rho)$ is isomorphic to $V \otimes B = \{(V \otimes B)_\alpha = V_1 \otimes B_\alpha\}_{\alpha \in \pi}$ as right B-module.*

Proof. Firstly, we‘ll prove that $L_{1,\alpha}\eta_\alpha g_1 = (I_1 \otimes \eta_\alpha g_1)\Delta^C_{1,1}$.

$$\begin{aligned}
&L_{1,\alpha}\eta_\alpha g_1\\
&= (\sigma_1 \otimes I^H_\alpha)\Delta^H_{1,\alpha}\eta_\alpha g_1\\
&= (\sigma_1 \otimes I^H_\alpha)(I_1 \otimes \eta_\alpha)\Delta^H_{1,1} g_1\\
&= (I_1 \otimes \eta_\alpha)(\sigma_1 \otimes I^H_1)\Delta^H_{1,1} g_1\\
&= (I_1 \otimes \eta_\alpha)L_{1,1} g_1\\
&= (I_1 \otimes \eta_\alpha)(I_1 \otimes g_1)\Delta^C_{1,1}\\
&= (I_1 \otimes \eta_\alpha g_1)\Delta^C_{1,1}.
\end{aligned}$$

We define $T_\alpha = (I_1 \otimes \eta_\alpha g_1)\rho_{1,1} : V_1 \to Ind(\rho)_\alpha$. We‘ll prove $T_\alpha(v_1) \in Ind(\rho)_\alpha$.

$$\begin{aligned}
&(I_1 \otimes L_{1,\alpha})T_\alpha(v_1)\\
&= (I_1 \otimes L_{1,\alpha})(I_1 \otimes \eta_\alpha g_1)\rho_{1,1}(v_1)\\
&= (I_1 \otimes L_{1,\alpha}\eta_\alpha g_1)\rho_{1,1}(v_1)\\
&= (I_1 \otimes (I_1 \otimes \eta_\alpha g_1)\Delta^C_{1,1})\rho_{1,1}(v_1)\\
&= (I_1 \otimes I_1 \otimes \eta_\alpha g_1)(I_1 \otimes \Delta^C_{1,1})\rho_{1,1}(v_1)\\
&= (I_1 \otimes I_1 \otimes \eta_\alpha g_1)(\rho_{1,1} \otimes I_1)\rho_{1,1}(v_1)\\
&= (\rho_{1,1} \otimes I_\alpha)(I_1 \otimes \eta_\alpha g_1)\rho_{1,1}(v_1)\\
&= (\rho_{1,1} \otimes I_\alpha)T_\alpha(v_1).
\end{aligned}$$

For $\alpha \in \pi$ we define $q_\alpha : V_1 \otimes B_\alpha \to Ind(\rho)_\alpha$ where

$$q_\alpha(v \otimes b) = \lambda_\alpha(T_\alpha(v) \otimes b) = v_1 \otimes \eta_\alpha g_1(v_2)b.$$

Clear $q_\alpha(v \otimes b) \in Ind(\rho)_\alpha$ and q_α is linear. We define $q_\alpha^{-1} : Ind(\rho)_\alpha \to V_1 \otimes B_\alpha$;

$$\begin{aligned}
q_\alpha^{-1} &= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(\rho_{1,1} \otimes I_\alpha) && \text{(i)}\\
&= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(I_1 \otimes L_{1,\alpha}) && \text{(ii)}
\end{aligned}$$

$$i.e., q_\alpha^{-1}(v \otimes h) = v_1 \otimes \eta_\alpha g_1^{-1}(v_2)h = v \otimes \eta_\alpha g_1^{-1}\sigma_1(h^1_1)h^\alpha_2.$$

We‘ll prove that $q_\alpha^{-1}(v\otimes h)\in V_1\otimes B_\alpha$.

$$\begin{aligned}
&(I_1\otimes L_{1,\alpha})q_\alpha^{-1}(v\otimes h)\\
&=(I_1\otimes L_{1,\alpha})(I_1\otimes\mu_\alpha(\eta_\alpha g_1^{-1}\otimes I_\alpha))(I_1\otimes L_{1,\alpha})(v\otimes h) \qquad \text{by equation (ii)}\\
&=(I_1\otimes\mu_{C_1\otimes H_\alpha}(L_{1,\alpha}\eta_\alpha g_1^{-1}\otimes L_{1,\alpha}))(I_1\otimes L_{1,\alpha})(v\otimes h)\\
&=v\otimes\mu_{C_1\otimes H_\alpha}((I_1\otimes\eta_\alpha)L_{1,1}g_1^{-1}\otimes L_{1,\alpha})L_{1,\alpha}(h)\\
&=v\otimes\mu_{C_1\otimes H_\alpha}((I_1\otimes\eta_\alpha)(S_1^C\otimes g_1^{-1})\tau\Delta^H_{1,1}\otimes L_{1,\alpha})L_{1,\alpha}(h) \qquad \text{by lemma 3.5}\\
&=v\otimes\mu_{C_1\otimes H_\alpha}((S_1^C\otimes\eta_\alpha g_1^{-1})\tau\Delta^H_{1,1}\otimes(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha})(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}(h)\\
&=v\otimes\mu_{C_1\otimes H_\alpha}((S_1^C\otimes\eta_\alpha g_1^{-1})\tau\otimes\sigma_1\otimes I^H_\alpha)(\Delta^H_{1,1}\otimes\Delta^H_{1,\alpha})(\sigma_1\otimes I^H_\alpha)\Delta^H_{1,\alpha}(h)\\
&=v\otimes\mu_{C_1\otimes H_\alpha}((S_1^C\otimes\eta_\alpha g_1^{-1})\tau\otimes\sigma_1\otimes I^H_\alpha)(\sigma_1\otimes\sigma_1\otimes I^H_1\otimes I^H_\alpha)(\Delta^H_{1,1}\otimes\Delta^H_{1,\alpha})\Delta^H_{1,\alpha}(h)\\
&=v\otimes S_1^C\sigma_1(h^1_{121})\sigma_1(h^1_{122})\otimes\eta_\alpha g_1^{-1}\sigma_1(h^1_{11}))h^\alpha_2\\
&=v\otimes\epsilon^C\sigma_1(h^1_{12})1\otimes\eta_\alpha g_\alpha^{-1}\sigma_1(h^1_{11}))h^\alpha_2\\
&=v\otimes\epsilon^H(h^1_{12})1\otimes\eta_\alpha g_1^{-1}\sigma_1(h^1_{11}))h^\alpha_2\\
&=v\otimes 1\otimes\eta_\alpha g_1^{-1}\sigma_1(\epsilon^H(h^1_{12})h^1_{11}))h^\alpha_2\\
&=v\otimes 1\otimes\eta_\alpha g_1^{-1}\sigma_1(h^1_1)h^\alpha_2\\
&=v\otimes 1\otimes\mu_\alpha(\eta_\alpha g_1^{-1}\otimes I_\alpha))L_{1,\alpha}(h)
\end{aligned}$$

Now, we prove that $q_\alpha q_\alpha^{-1}=I$ and $q_\alpha^{-1}q_\alpha=I$.

$$\begin{aligned}
&q_\alpha q_\alpha^{-1}(v\otimes h)\\
&=(I_1\otimes\mu_\alpha)(I_1\otimes\eta_\alpha g_1\otimes I_\alpha)(\rho_{1,1}\otimes I_\alpha)\\
&\quad(I_1\otimes\mu_\alpha(\eta_\alpha g_1^{-1}\otimes I_\alpha))(\rho_{1,1}\otimes I_\alpha)(v\otimes h) \qquad \text{by equation (ii)}\\
&=(I_1\otimes\mu_\alpha)(I_1\otimes\eta_\alpha g_1\otimes I_\alpha)(I_1\otimes I_1\otimes\mu_\alpha(\eta_\alpha g_1^{-1}\otimes I_\alpha))(\rho_{1,1}\otimes I_1\otimes I_\alpha)(\rho_{1,1}\otimes I_\alpha)(v\otimes h)\\
&=(I_1\otimes\mu_\alpha)(I_1\otimes\eta_\alpha g_1\otimes I_\alpha)(I_1\otimes I_1\otimes\mu_\alpha(\eta_\alpha g_1^{-1}\otimes I_\alpha))((\rho_{1,1}\otimes I_1)\rho_{1,1}\otimes I_\alpha)(v\otimes h)\\
&=(I_1\otimes\mu_\alpha)(I_1\otimes\eta_\alpha g_1\otimes I_\alpha)(I_1\otimes I_1\otimes\mu_\alpha(\eta_\alpha g_1^{-1}\otimes I_\alpha))((I_1\otimes\Delta^C_{1,1}\otimes I_\alpha)(\rho_{1,1}\otimes I_\alpha)(v\otimes h)\\
&=v^1_1\otimes\eta_\alpha g_1(v^1_{21})\eta_\alpha g_1^{-1}(v^1_{22})h\\
&=v^1_1\otimes\eta_\alpha(g_1(v^1_{21})g_1^{-1}(v^1_{22}))h\\
&=v^1_1\otimes\eta_\alpha(\epsilon(v^1_2)1)h\\
&=\epsilon(v^1_2)v^1_1\otimes\eta_\alpha(1)h\\
&=v\otimes h
\end{aligned}$$

$$\begin{aligned}
& q_\alpha^{-1} q_\alpha(v \otimes b) \\
&= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(I_1 \otimes L_{1,\alpha})(I_1 \otimes \mu_\alpha)(I_1 \otimes \eta_\alpha g_1 \otimes I_\alpha)(\rho_{1,1} \otimes I_\alpha)(v \otimes b) \text{by equation (i)} \\
&= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(I_1 \otimes L_{1,\alpha})(v_1 \otimes \eta_\alpha g_1(v_2^1)b) \\
&= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(v_1 \otimes L_{1,\alpha}(\eta_\alpha g_1(v_2^1))L_{1,\alpha}(b)) \\
&= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(v_1 \otimes (I_1 \otimes \eta_\alpha)L_{1,1} g_1(v_2^1) \cdot (1 \otimes b)) \\
&= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(v_1 \otimes (I_1 \otimes \eta_\alpha)(I_1 \otimes g_1)\Delta_{1,1}^C(v_2^1) \cdot (1 \otimes b)) \\
&= (I_1 \otimes \mu_\alpha(\eta_\alpha g_1^{-1} \otimes I_\alpha))(v_1 \otimes v_{21}^1 \otimes \eta_\alpha g_1(v_{22}^1)b) \\
&= v_1 \otimes \eta_\alpha g_1^{-1}(v_{21}^1)\eta_\alpha g_1(v_{22}^1)b \\
&= v_1 \otimes \eta_\alpha(g_1^{-1}(v_{21}^1)g_1(v_{22}^1))b \\
&= v_1 \otimes \eta_\alpha(\epsilon^C(v_2^1)1)b \\
&= \epsilon^C(v_2^1)v_1 \otimes \eta_\alpha(1)b \\
&= v \otimes b.
\end{aligned}$$

Now we prove that q_α is module map for all $\alpha \in \pi$,

$$\begin{aligned}
& q_\alpha(I_1 \otimes \mu_\alpha)(v \otimes b \otimes k) \\
&= q_\alpha(v \otimes bk) \\
&= \lambda_\alpha(T_\alpha(v) \otimes bk) \\
&= \lambda_\alpha((I_1 \otimes \eta_\alpha g_1)\rho_{1,1}(v) \otimes bk) \\
&= \lambda_\alpha(v_1 \otimes \eta_\alpha g_1(v_2) \otimes bk) \\
&= v_1 \otimes \eta_\alpha g_1(v_2)bk \\
&= v_1 \otimes (\eta_\alpha g_1(v_2)b)k \\
&= \lambda_\alpha(v_1 \otimes (\eta_\alpha g_1(v_2)b) \otimes k) \\
&= \lambda_\alpha(q_\alpha \otimes I_\alpha)(v \otimes b \otimes k).
\end{aligned}$$

∎

Throughout this last part, we consider (C, σ) as a left $\pi-$ coisotropic quantum subgroup of H, $V = \{V_\alpha\}_{\alpha\in\pi}$ as a right $\pi-$comodule over C and $G = \{G_\alpha\}_{\alpha\in\pi}$,where $G_\alpha = \{h \in H_\alpha : L_{1,\alpha}(h) = (\sigma_1 \otimes I_\alpha^H)\Delta_{1,\alpha}^H(h) = \sigma_1(1) \otimes h\}$.

Lemma 3.13. *$Ind(\rho)$ is right G-module*

Proof. Similar to lemma 3.1. ∎

Theorem 3.14. *H is isomorphic to $C \otimes G$ as vector space.*

Proof. 1. For $h \in H_\alpha$, as in lemma 3.6, $h = g_\alpha \sigma_\alpha(h_1^\alpha) g_\alpha^{-1} \sigma_{\alpha^{-1}}(h_{21}^{\alpha^{-1}}) h_{22}^\alpha$.

2. For $h \in H_1$ and $\alpha \in \pi$, we'll prove that $\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h) \in G_\alpha$.

$$\begin{aligned}
& L_{1,\alpha}\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (\sigma_1 \otimes I^H_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (\sigma_1 \otimes I^H_\alpha)\mu_{H_1\otimes H_\alpha}(\Delta^H_{1,\alpha} \otimes \Delta^H_{1,\alpha})(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (\sigma_1\mu_{H_1} \otimes \mu_{H_\alpha})(I \otimes \tau \otimes I)(\Delta^H_{1,\alpha} \otimes \Delta^H_{1,\alpha})(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I^H_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (I_1 \otimes \mu_{H_\alpha})((\sigma_1\mu_{H_1} \otimes I_\alpha)(I \otimes \tau) \otimes I^H_\alpha) \\
& (\Delta^H_{1,\alpha}g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_1 \otimes I^H_\alpha)(I_{\alpha^{-1}} \otimes \Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (I_1 \otimes \mu_{H_\alpha})((\sigma_1\mu_{H_1} \otimes I_\alpha)(I \otimes \tau)(\Delta^H_{1,\alpha}g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_1) \otimes I^H_\alpha)(I_{\alpha^{-1}} \otimes \Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (I_1 \otimes \mu_{H_\alpha})((I_1 \otimes g_\alpha^{-1})(\sigma_1 \otimes \sigma_{\alpha^{-1}})(\mu_1(S^H_1 \otimes I_1) \otimes I_{\alpha^{-1}}) \\
& (I_1 \otimes \tau)(\tau\Delta^H_{\alpha^{-1},1} \otimes I_1) \otimes I^H_\alpha)(I_{\alpha^{-1}} \otimes \Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (I_1 \otimes \mu_{H_\alpha})((I_1 \otimes g_\alpha^{-1})(\sigma_1 \otimes \sigma_{\alpha^{-1}})(\mu_1(S^H_1 \otimes I_1) \otimes I_{\alpha^{-1}}) \\
& (I_1 \otimes \tau)(\tau \otimes I_1) \otimes I^H_\alpha)(\Delta^H_{\alpha^{-1},1} \otimes \Delta^H_{1,\alpha})\Delta^H_{\alpha^{-1},\alpha}(h) \\
= {} & (I_1 \otimes \mu_{H_\alpha})[(\sigma_1 \otimes g_\alpha^{-1}\sigma_{\alpha^{-1}})(\mu_1(S^H_1 \otimes I_1) \otimes I_{\alpha^{-1}}) \\
& (I_1 \otimes \tau)(\tau \otimes I_1) \otimes I^H_\alpha](h^{\alpha^{-1}}_{11} \otimes h^1_{121} \otimes h^1_{122} \otimes h^\alpha_2) \\
= {} & \sigma_1(S^H_1(h^1_{121})h^1_{122}) \otimes g_\alpha^{-1}(\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{11}))h^\alpha_2 \\
= {} & \sigma_1(\epsilon^H(h^1_{12})1) \otimes g_\alpha^{-1}(\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{11}))h^\alpha_2 \\
= {} & \sigma_1(1) \otimes g_\alpha^{-1}(\sigma_{\alpha^{-1}}(\epsilon^H(h^1_{12})h^{\alpha^{-1}}_{11}))h^\alpha_2 \\
= {} & \sigma_1(1) \otimes g_\alpha^{-1}(\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_1))h^\alpha_2 \\
= {} & \sigma_1(1) \otimes \mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha)\Delta^H_{\alpha^{-1},\alpha}(h)
\end{aligned}$$

3. We define $A_\alpha : C_\alpha \otimes G_\alpha \to H_\alpha$ as follow

$$A_\alpha(c \otimes b) = \mu_\alpha(g_\alpha \otimes I_\alpha)(c \otimes b) = g_\alpha(c)b.$$

Clear A_α is linear. We define $A_\alpha^{-1} : H_\alpha \to C_\alpha \otimes G_\alpha$ as follow

$$A_\alpha^{-1}(h) = (\sigma_\alpha \otimes \mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha))(\Delta^H_{\alpha,\alpha^{-1}} \otimes I_\alpha)\Delta^H_{1,\alpha}(h).$$

By step 2 and property of Δ , we have $A_\alpha^{-1}(h) \in C_\alpha \otimes G_\alpha$. We will prove $A_\alpha A_\alpha^{-1} = I_{H_\alpha}$ and $A_\alpha^{-1}A_\alpha = I_{C_\alpha \otimes B_\alpha}$. Let $h \in H_\alpha, c \otimes b \in C_\alpha \otimes G_\alpha$,

$$\begin{aligned}
& A_\alpha A_\alpha^{-1}(h) \\
= {} & A_\alpha((\sigma_\alpha \otimes \mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha))(\Delta^H_{\alpha,\alpha^{-1}} \otimes I_\alpha)\Delta^H_{1,\alpha}(h)) \\
= {} & A_\alpha((\sigma_\alpha \otimes \mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}} \otimes I_\alpha))(I_\alpha \otimes \Delta^H_{\alpha^{-1},\alpha})\Delta^H_{\alpha,1}(h)) \\
= {} & A_\alpha(\sigma_\alpha(h^\alpha_1) \otimes g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{21})h^\alpha_{22}) \\
= {} & g_\alpha\sigma_\alpha(h^\alpha_1)g_\alpha^{-1}\sigma_{\alpha^{-1}}(h^{\alpha^{-1}}_{21})h^\alpha_{22} \\
= {} & h.
\end{aligned}$$

and

$$\begin{aligned}
&A_\alpha^{-1}A_\alpha(c\otimes k)\\
&=(\sigma_\alpha\otimes\mu_\alpha(g_\alpha^{-1}\sigma_{\alpha^{-1}}\otimes I_\alpha))(\Delta^H_{\alpha,\alpha^{-1}}\otimes I_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha\otimes I_\alpha)(c\otimes k)\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\sigma_\alpha\otimes\sigma_{\alpha^{-1}}\otimes I^H_\alpha)(\Delta^H_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha\otimes I_\alpha)(c\otimes k)\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)((\sigma_\alpha\otimes\sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)\Delta^H_{1,\alpha}\mu_\alpha(g_\alpha\otimes I_\alpha)(c\otimes k)\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\sigma_1\otimes I^H_\alpha)\mu_{H_\alpha\otimes H_{\alpha^{-1}}}(\Delta^H_{1,\alpha}g_\alpha\otimes\Delta^H_{1,\alpha})(c\otimes k)\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes I^H_\alpha)(\sigma_1\mu_1\otimes\mu_\alpha)(I\otimes\tau\otimes I)(\Delta^H_{1,\alpha}g_\alpha\otimes\Delta^H_{1,\alpha})(c\otimes k)\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes\mu_\alpha)[(\sigma_1\mu_1\otimes I^H_\alpha)(I\otimes\tau)(\Delta^H_{1,\alpha}g_\alpha(c)\otimes k^1_1\otimes k^\alpha_2)]\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes\mu_\alpha)\\
&\quad[(I_1\otimes g_\alpha)(\sigma_1\otimes\sigma_\alpha)(\mu_1\otimes I_\alpha)(I_1\otimes\tau)(\Delta^C_{1,\alpha}\otimes I_1)(v\otimes k^1_1)\otimes k^\alpha_2]\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes\mu_\alpha)[\sigma_1(v^1_1k^1_1)\otimes g_\alpha\sigma_\alpha(v^\alpha_2)\otimes k^\alpha_2]\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes\mu_\alpha)[\omega_1(v^1_1\otimes\sigma_1(k^1_1))\otimes g_\alpha\sigma_\alpha(v^\alpha_2)\otimes k^\alpha_2]\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes\mu_\alpha)[\omega_1(v^1_1\otimes\sigma_1(1))\otimes g_\alpha\sigma_\alpha(v^\alpha_2)\otimes k]\\
&=(I_\alpha\otimes\mu_\alpha)(I_\alpha\otimes g_\alpha^{-1}\otimes I_\alpha)(\Delta^C_{\alpha,\alpha^{-1}}\otimes\mu_\alpha)[\sigma_1(v^1_1)\otimes g_\alpha\sigma_\alpha(v^\alpha_2)\otimes k]\\
&=(I_\alpha\otimes\mu_\alpha)[(I_\alpha\otimes g_\alpha^{-1})\Delta^C_{\alpha,\alpha^{-1}}\sigma_1(v^1_1)\otimes g_\alpha\sigma_\alpha(v^\alpha_2)k]\\
&=(I_\alpha\otimes\mu_\alpha)[(I_\alpha\otimes g_\alpha^{-1})(\sigma_\alpha\otimes\sigma_{\alpha^{-1}})\Delta^H_{\alpha,\alpha^{-1}}(v^1_1)\otimes g_\alpha\sigma_\alpha(v^\alpha_2)k]\\
&=(I_\alpha\otimes\mu_\alpha)[(I_\alpha\otimes g_\alpha^{-1})(\sigma_\alpha(v^\alpha_1)\otimes\sigma_{\alpha^{-1}}(v^{\alpha^{-1}}_2)\otimes g_\alpha\sigma_\alpha(v^\alpha_3)k]\\
&=\sigma_\alpha(v^\alpha_1)\otimes g_\alpha^{-1}\sigma_{\alpha^{-1}}(v^{\alpha^{-1}}_2)g_\alpha\sigma_\alpha(v^\alpha_3)k\\
&=\sigma_\alpha(v^\alpha_1)\otimes\epsilon^C\sigma_1(v^1_2)k\\
&=\sigma_\alpha(\epsilon^H(v^1_2)v^\alpha_1)\otimes k\\
&=\sigma_\alpha(v)\otimes k\\
&=c\otimes k.
\end{aligned}$$

∎

4 Coinduced representations of Hopf group coalgebra

In this section, we study coinduced representation from left $\pi-$coisotropic quantum subgroup. We restrict our attention to finite dimensional case of Hopf $\pi-$coalgebra, i.e., dim $H_\alpha = n^\alpha \prec \infty$ for all $\alpha \in \pi$. Let us start now from left $\pi-$corepresentation $\rho = \{\rho_{\alpha,\beta}\}_{\alpha,\beta\pi}$ from left $\pi-$coisotropic quantum subgroup (C, σ) on $V = \{V_\alpha\}_{\alpha\in\pi}$. We define $W = \{W_\alpha\}_{\alpha\in\pi}$, where $W_\alpha = \{F_\alpha \in Hom(V_1, H_\alpha) :\ L_{1,\alpha}F_\alpha = (I_{C_1} \otimes F_\alpha)\rho_{1,1}\}$ and $L_{\alpha,\beta} = (\sigma_\alpha \otimes I_\beta)\Delta_{\alpha,\beta}$

Lemma 4.1. *For* $\alpha, \beta, \gamma \in \pi$, $(L_{\alpha,\beta} \otimes I_\gamma)\Delta_{\alpha\beta,\gamma} = (I_\alpha \otimes \Delta_{\beta,\gamma})L_{\alpha,\beta\gamma}$

Proof. Similar to lemma 2.4 . ∎

Lemma 4.2. *Suppose that* $\{e_i^\alpha\}_{i=1}^{n^\alpha}$ *and* $\{g_i^\alpha\}_{i=1}^{n^\alpha}$ *are dual bases of* H_α *and* H_α^* *for all* $\alpha \in \pi$*. For* $\alpha, \beta, \gamma \in \pi$*, fix* $F_{\alpha\beta\gamma} \in W_{\alpha\beta\gamma}$*, if we define the two maps* $\xi_1 : V_1 \to H_\alpha \otimes H_\beta \otimes H_\gamma$ *and* $\xi_2 : V_1 \to H_\alpha \otimes H_\beta \otimes H_\gamma$ *such that*

$$\xi_1(v_1) = \sum_{i=1}^{n^{\beta\gamma}} \sim (I_\alpha \otimes g_i^{\beta\gamma})\Delta_{\alpha,\beta\gamma} F_{\alpha\beta\gamma}(v_1) \otimes \Delta_{\beta,\gamma}(e_i^{\beta\gamma})$$

$$\xi_2(v_1) = \sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma} [\sim (\sim \otimes I_K)(I_\alpha \otimes (g_h^\beta \otimes g_l^\gamma)\Delta_{\beta,\gamma}]\Delta_{\alpha,\beta\gamma} F_{\alpha\beta\gamma}(v_1) \otimes e_h^\beta \otimes e_l^\gamma$$

then $\xi_1 = \xi_2$*.*

Proof. We put $\quad F_{\alpha\beta\gamma}(v_1) = h_{\alpha\beta\gamma} = \sum_{j=1}^{n^{\alpha\beta\gamma}} \lambda_j e_j^{\alpha\beta\gamma}$
$\Delta_{\beta,\gamma}(e_i^{\beta\gamma}) = \sum_{r=1}^{n^\beta}\sum_{s=1}^{n^\gamma} \eta_{rs}^i e_r^\beta \otimes e_s^\gamma$ and $\Delta_{\alpha,\beta\gamma}(e_i^{\alpha\beta\gamma}) = \sum_{r=1}^{n^\alpha}\sum_{s=1}^{n^{\beta\gamma}} \theta_{rs}^i e_r^\alpha \otimes e_s^{\beta\gamma}$
then we have

$$\Delta_{\alpha,\beta\gamma} F_{\alpha\beta\gamma}(v_1) = \Delta_{\alpha,\beta\gamma}(h_{\alpha\beta\gamma}) = \sum_{j=1}^{n^\gamma} \lambda_j \Delta_{\alpha,\beta\gamma}(e_j^{\alpha\beta\gamma}) = \sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{s=1}^{n^{\beta\gamma}} \lambda_j \theta_{rs}^j e_r^\alpha \otimes e_s^{\beta\gamma}$$

imply that

$$\begin{aligned}
&\sum_{i=1}^{n^{\beta\gamma}} \sim (I_\alpha \otimes g_i^{\beta\gamma})\Delta_{\alpha,\beta\gamma} F_{\alpha\beta\gamma}(v_1) \otimes \Delta_{\beta,\gamma}(e_i^{\beta\gamma}) \\
&\quad = \sum_{i=1}^{n^{\beta\gamma}} [\sim (I_\alpha \otimes g_i^{\beta\gamma}) \sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{s=1}^{n^{\beta\gamma}} \lambda_j \theta_{rs}^j e_r^\alpha \otimes e_s^{\beta\gamma}] \otimes \sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma} \eta_{hl}^i e_h^\beta \otimes e_l^\gamma \\
&\quad = \sum_{i=1}^{n^{\beta\gamma}}\sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{s=1}^{n^{\beta\gamma}} \lambda_j \theta_{rs}^j g_i^{\beta\gamma}(e_s^{\beta\gamma}) e_r^\alpha \otimes \sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma} \eta_{hl}^i e_h^\beta \otimes e_l^\gamma \\
&\quad = \sum_{i=1}^{n^{\beta\gamma}}\sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha} \lambda_j \theta_{ri}^j e_r^\alpha \otimes \sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma} \eta_{hl}^i e_h^\beta \otimes e_l^\gamma \\
&\quad = \sum_{i=1}^{n^{\beta\gamma}}\sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma} \eta_{hl}^i \lambda_j \theta_{ri}^j e_r^\alpha \otimes e_h^\beta \otimes e_l^\gamma.
\end{aligned}$$

$$
\begin{aligned}
&\sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma}[\sim(\sim\otimes I_K)(I_\alpha\otimes(g_h^\beta\otimes g_l^\gamma)\Delta_{\beta,\gamma}]\Delta_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}(v_1)\otimes e_h^\beta\otimes e_l^\gamma\\
&\qquad=\sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma}[\sim(\sim\otimes I_K)(I_\alpha\otimes(g_h^\beta\otimes g_l^\gamma)\Delta_{\beta,\gamma}][\sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{i=1}^{n^{\beta\gamma}}\lambda_j\theta_{ri}^j e_r^\alpha\otimes e_i^{\beta\gamma}]\otimes e_h^\beta\otimes e_l^\gamma\\
&\qquad=\sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma}\sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{i=1}^{n^{\beta\gamma}}\lambda_j\theta_{ri}^j\sim(\sim\otimes I_K)[e_r^\alpha\otimes(g_h^\beta\otimes g_l^\gamma)\Delta_{\beta,\gamma}(e_i^{\beta\gamma})]\otimes e_h^\beta\otimes e_l^\gamma\\
&\qquad=\sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma}\sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{i=1}^{n^{\beta\gamma}}\sum_{q=1}^{n^\beta}\sum_{s=1}^{n^\gamma}\lambda_j\theta_{ri}^j\eta_{qs}^i g_h^\beta(e_q^\beta)g_l^\gamma(e_s^\gamma)e_r^\alpha\otimes e_h^\beta\otimes e_l^\gamma\\
&\qquad=\sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma}\sum_{j=1}^{n^\gamma}\sum_{r=1}^{n^\alpha}\sum_{i=1}^{n^{\beta\gamma}}\eta_{hl}^i\lambda_j\theta_{ri}^j e_r^\alpha\otimes e_h^\beta\otimes e_l^\gamma
\end{aligned}
$$

Therefore, we have $\sum_{i=1}^{n^{\beta\gamma}}\sim(I_\alpha\otimes g_i^{\beta\gamma})\Delta_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}(v_1)\otimes\Delta_{\beta,\gamma}(e_i^{\beta\gamma})=\sum_{h=1}^{n^\beta}\sum_{l=1}^{n^\gamma}[\sim(\sim\otimes I_K)(I_\alpha\otimes(g_h^\beta\otimes g_l^\gamma)\Delta_{\beta,\gamma}]\Delta_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}(v_1)\otimes e_h^\beta\otimes e_l^\gamma$, and hence $\xi_1=\xi_2$ ∎

Theorem 4.3. $W=\{W_\alpha\}_{\alpha\in\pi}$ *is right* $\pi-$*comodule over* H *by* $\Omega=\{\Omega_{\alpha,\beta}\}_{\alpha,\beta\in\pi}$ *as*

$$\Omega_{\alpha,\beta}:W_{\alpha\beta}\to W_\alpha\otimes H_\beta \text{ as } \Omega_{\alpha,\beta}(F_{\alpha\beta})=\sim(I_\alpha\otimes g^\beta)\Delta_{\alpha,\beta}F_{\alpha\beta}\otimes e^\beta.$$

Proof. Firstly, we 'll prove that $\sim(I_\alpha\otimes g^\beta)\Delta_{\alpha,\beta}F_{\alpha\beta}\in W_\alpha$.

$$
\begin{aligned}
&L_{1,\alpha}[\sim(I_\alpha\otimes g^\beta)\Delta_{\alpha,\beta}F_{\alpha\beta}]\\
&=\sim(L_{1,\alpha}\otimes g^\beta)\Delta_{\alpha,\beta}F_{\alpha\beta}\\
&=(I_{C_1}\otimes\sim(I_\alpha\otimes g^\beta))(L_{1,\alpha}\otimes I_\beta)\Delta_{\alpha,\beta}F_{\alpha\beta}\\
&=(I_{C_1}\otimes\sim(I_\alpha\otimes g^\beta))(I_{C_1}\otimes\Delta_{\alpha,\beta})L_{1,\alpha\beta}F_{\alpha\beta}\\
&=(I_{C_1}\otimes\sim(I_\alpha\otimes g^\beta))(I_{C_1}\otimes\Delta_{\alpha,\beta})(I_{C_1}\otimes F_{\alpha\beta})\rho_{1,1}\\
&=(I_{C_1}\otimes\sim(I_\alpha\otimes g^\beta)\Delta_{\alpha,\beta}F_{\alpha\beta})\rho_{1,1}
\end{aligned}
$$

Now, we will prove that Ω is coaction on W, i.e., the two diagrams

$$\begin{tikzcd}
W_{\alpha\beta\gamma} \arrow[r, "\Omega_{\alpha\beta,\gamma}"] \arrow[d, "\Omega_{\alpha,\beta\gamma}"'] & W_{\alpha\beta}\otimes H_\gamma \arrow[d, "(\Omega_{\alpha,\beta}\otimes I_\gamma)"] \\
W_\alpha\otimes H_{\beta\gamma} \arrow[r, "I_{W_\alpha}\otimes\Delta_{\beta,\gamma}"'] & W_\alpha\otimes H_\beta\otimes H_\gamma
\end{tikzcd}
,\qquad
\begin{tikzcd}
W_\alpha \arrow[r, "\Omega_{\alpha,1}"] \arrow[dr, "\sim"'] & W_\alpha\otimes H_1 \arrow[d, "(I_{W_\alpha}\otimes\epsilon)"] \\
& W_\alpha\otimes K
\end{tikzcd}$$

are commutes.

$$\begin{aligned}
&(\Omega_{\alpha,\beta}\otimes I_\gamma)\Omega_{\alpha\beta,\gamma}F_{\alpha\beta\gamma}\\
&=(\Omega_{\alpha,\beta}\otimes I_\gamma)(\sim(I_{\alpha\beta}\otimes g^\gamma)\Delta_{\alpha\beta,\gamma}F_{\alpha\beta\gamma}\otimes e^\gamma)\\
&=\Omega_{\alpha,\beta}(\sim(I_{\alpha\beta}\otimes g^\gamma)\Delta_{\alpha\beta,\gamma}F_{\alpha\beta\gamma})\otimes e^\gamma\\
&=\sim(I_\alpha\otimes g^\beta)\Delta_{\alpha,\beta}\sim(I_{\alpha\beta}\otimes g^\gamma)\Delta_{\alpha\beta,\gamma}F_{\alpha\beta\gamma}\otimes e^\beta\otimes e^\gamma\\
&=\sim(\sim\otimes I_K)((I_\alpha\otimes g^\beta)\Delta_{\alpha,\beta}\otimes g^\gamma)\Delta_{\alpha\beta,\gamma}F_{\alpha\beta\gamma}\otimes e^\beta\otimes e^\gamma\\
&=\sim(\sim\otimes I_K)(I_\alpha\otimes g^\beta\otimes g^\gamma)(\Delta_{\alpha,\beta}\otimes I_\gamma)\Delta_{\alpha\beta,\gamma}F_{\alpha\beta\gamma}\otimes e^\beta\otimes e^\gamma\\
&=\sim(\sim\otimes I_K)(I_\alpha\otimes g^\beta\otimes g^\gamma)(I_\alpha\otimes\Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}\otimes e^\beta\otimes e^\gamma\\
&=\sim(\sim\otimes I_K)(I_\alpha\otimes(g^\beta\otimes g^\gamma)\Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}\otimes e^\beta\otimes e^\gamma\\
&=\xi_1
\end{aligned}$$

Also

$$\begin{aligned}
&(I_{W_\alpha}\otimes\Delta_{\beta,\gamma})\Omega_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}\\
&=(I_{W_\alpha}\otimes\Delta_{\beta,\gamma})(\sim(I_\alpha\otimes g^{\beta\gamma})\Delta_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}\otimes e^{\beta\gamma})\\
&=\sim(I_{W_\alpha}\otimes g^{\beta\gamma})\Delta_{\alpha,\beta\gamma}F_{\alpha\beta\gamma}\otimes\Delta_{\beta,\gamma}(e^{\beta\gamma})\\
&=\xi_2.
\end{aligned}$$

from lemma 4.2. the first diagram is commute. For the second diagram

$$\begin{aligned}
&(I_{W_\alpha}\otimes\epsilon)\Omega_{\alpha,1}F_\alpha\\
&=(I_{W_\alpha}\otimes\epsilon)[\sim(I_\alpha\otimes g^1)\Delta_{\alpha,1}F_\alpha\otimes e^1]\\
&=\sim(I_{W_\alpha}\otimes g^1)\Delta_{\alpha,1}F_\alpha\otimes\epsilon(e^1)\\
&=\sim(I_{W_\alpha}\otimes\epsilon(e^1)g^1)\Delta_{\alpha,1}F_\alpha\otimes 1_K\\
&=\sim(I_{W_\alpha}\otimes\epsilon)\Delta_{\alpha,1}F_\alpha\otimes 1_K\\
&=F_\alpha\otimes 1_K.
\end{aligned}$$

▮

Remark 4.4. Given a right corepresentation ρ of the $\pi-$coisotropic quantum subgroup of (C,σ) the corresponding corepresentation $(I\otimes\Delta)$ on $Coind(\rho)$ of H is called coinduced representation from ρ on H.

Now, we constract an induced and coinduced representation from subHopf $\pi-$coalgebra.

Theorem 4.5. *Let H be a finite dimentional Hopf π-coalgebra and $A=\{A_\alpha\}_{\alpha\in\pi}$ be an isolated subHopf π-coalgebra of H. If $V=\{V_\alpha\}_{\alpha\in\pi}$ is left π-comodule over A by $\rho=\{\rho_{\alpha,\beta}\}_{\alpha,\beta\in\pi}$, then we can construct an induced and coinduced representation over H.*

Proof. Since $A = \{A_\alpha\}_{\alpha\in\pi}$ is an isolated subHopf π-coalgebra of H, there exist a family $I = \{I_\alpha\}_{\alpha\in\pi}$ of Hopf π-coideal of H such that $H_\alpha = A_\alpha \oplus I_\alpha$ for all $\alpha \in \pi$. We will prove that (A, σ) is left $\pi-$coisotropic quantum subgroup of H where $\sigma = \{\sigma_\alpha\}_{\alpha\in\pi}$ and $\sigma_\alpha : H_\alpha \to A_\alpha$ as $\sigma_\alpha(m_\alpha + i_\alpha) = m_\alpha$. Clear, A is π-coalgebra. We 'll prove that A_α is left H_α-module for all $\alpha \in \pi$. We define $\Phi_\alpha : H_\alpha \otimes A_\alpha \to A_\alpha$ as follow $\Phi_\alpha((m_\alpha + i_\alpha) \otimes a_\alpha) = m_\alpha a_\alpha$. Then $\Phi_\alpha(\mu_\alpha \otimes I_\alpha)((m_\alpha + i_\alpha) \otimes (n_\alpha + j_\alpha) \otimes a_\alpha) = \Phi_\alpha((m_\alpha n_\alpha + m_\alpha j_\alpha + i_\alpha n_\alpha + i_\alpha j_\alpha) \otimes a_\alpha) = m_\alpha n_\alpha a_\alpha$ and $\Phi_\alpha(I_\alpha \otimes \Phi_\alpha)((m_\alpha + i_\alpha) \otimes (n_\alpha + j_\alpha) \otimes a_\alpha) = \Phi_\alpha((m_\alpha + i_\alpha) \otimes n_\alpha a_\alpha) = m_\alpha n_\alpha a_\alpha$ where $m_\alpha j_\alpha + i_\alpha n_\alpha + i_\alpha j_\alpha \in I_\alpha$. Also, we have $\Phi_\alpha(\eta_\alpha \otimes I_\alpha)(k \otimes a_\alpha) = \Phi_\alpha((\eta_\alpha(k) + 0) \otimes a_\alpha) = \eta_\alpha(k)a_\alpha = ka_\alpha$. Now, we will prove that σ_α is $\pi-$coalgebra map, $(\sigma_\alpha \otimes \sigma_\beta)\Delta_{\alpha,\beta}(h_{\alpha\beta}) = (\sigma_\alpha \otimes \sigma_\beta)\Delta_{\alpha,\beta}(m_{\alpha\beta} + i_{\alpha\beta}) = (\sigma_\alpha \otimes \sigma_\beta)(\Delta_{\alpha,\beta}(m_{\alpha\beta}) + \Delta_{\alpha,\beta}(i_{\alpha\beta})) = \Delta_{\alpha,\beta}(m_{\alpha\beta})$ and $\Delta_{\alpha,\beta}\sigma_{\alpha\beta}(h_{\alpha\beta}) = \Delta_{\alpha,\beta}\sigma_{\alpha\beta}(m_{\alpha\beta}+i_{\alpha\beta}) = \Delta_{\alpha,\beta}(m_{\alpha\beta})$. We will prove that σ_α is left module map, $\Phi_\alpha(I_\alpha\otimes\sigma_\alpha)(h_\alpha\otimes k_\alpha) = \Phi_\alpha(I_\alpha\otimes\sigma_\alpha)((m_\alpha+i_\alpha)\otimes(n_\alpha+j_\alpha)) = \Phi_\alpha((m_\alpha + i_\alpha) \otimes n_\alpha) = m_\alpha n_\alpha$ and $\sigma_\alpha\mu_\alpha(h_\alpha \otimes k_\alpha) = \sigma_\alpha(m_\alpha n_\alpha + m_\alpha j_\alpha + i_\alpha n_\alpha + i_\alpha j_\alpha) = m_\alpha n_\alpha$ Therefore (A, σ) is left $\pi-$coisotropic quantum subgroup of H. Since V is left $\pi-$comodule over A, then we can construct an induced and coinduced representation over H, by theorem 2.6 and theorem 4.3. ∎

5 Simplicity of induced representation of Hopf group coalgebra

In this section we have studied the simplicity of our induced representations of Hopf $\pi-$coalgebra. Let H be a Hopf $\pi-$coalgebra, (C, σ) a $\pi-$coisotropic quantum subgroup. If we have two equivalent right $\pi-$comodules over C. What is the relations between the two induced and the two coinduced representations over H ? Also, we have the following question: what is the algebraic relations of the induction with respect to direct sums of representations in point of view of simplicity. Finally, we have proved that if the induced representation is simple, then its algebraic induction is constructed from simple representation of the group coisotropic quantum subgroup. If we have simple representation of group coisotropic quantum subgroup for Hopf $\pi-$coalgebra, then the induced representation is not necessary to be simple.

A right π-comodule $M = \{M_\alpha\}_{\alpha\in\pi}$ is said to be simple if it is non-zero (i.e., $M_\alpha \neq 0$ for some $\alpha \in \pi$) and if it has no $\pi-$subcomodules other than $0 = 0_{\alpha\in\pi}$ and itself.

A right $\pi-$comodules (V, ρ) and (W, ρ') over C are equivalent if there exist a family $F = \{F_\alpha : V_\alpha \to W_\alpha\}_{\alpha\in\pi}$ of vector space isomorphisms such that

$$\rho'_{\alpha,\beta}F_{\alpha\beta} = (F_\alpha \otimes I_\beta)\rho_{\alpha,\beta} \quad \text{for all } \alpha, \beta \in \pi$$

Theorem 5.1. *Let (V, ρ) and (W, ρ') be a right $\pi-$comodules over a $\pi-$coisotropic quantum subgroup (C, σ). If there exist a vector space isomorphism F_1 : $V_1 \to W_1$ such that*

$$\psi_{1,1}F_1 = (F_1 \otimes I_1)\rho_{1,1} \tag{\#}$$

then the induced representations $Ind(\rho)$ and $Ind(\rho')$ are equivalent right $\pi-$comodules over H.

Proof. Let us consider the family of vector space isomorphisms

$$\overline{F} = \{\overline{F}_\alpha = F_1 \otimes I_\alpha : V_1 \otimes H_\alpha \to W_1 \otimes H_\alpha\}_{\alpha\in\pi}.$$

Firstly, we must prove that, for $\alpha \in \pi$, $\overline{F}_\alpha(Ind(\rho)_\alpha) \subseteq Ind(\rho')_\alpha$. Let $f \in Ind(\rho)_\alpha$, then

$$(I_1 \otimes L_{1,\alpha})(f) = (\rho_{1,1} \otimes I_\alpha)(f) \tag{1}$$

We want to prove $\overline{F}_\alpha(f) \in Ind(\rho')_\alpha$, i. e., $(I_1 \otimes L_{1,\alpha})\overline{F}_\alpha(f) = (\rho'_{1,1} \otimes I_\alpha)\overline{F}_\alpha(f)$

$$\begin{aligned}
&(I_1 \otimes L_{1,\alpha})\overline{F}_\alpha(f) \\
&= (I_1 \otimes L_{1,\alpha})(F_1 \otimes I_\alpha)(f) \\
&= (F_1 \otimes I_1 \otimes I_\alpha)(I_1 \otimes L_{1,\alpha})(f) \\
&= (F_1 \otimes I_1 \otimes I_\alpha)(\rho_{1,1} \otimes I_\alpha)(f) && \text{by (1)} \\
&= [(F_1 \otimes I_1)\rho_{1,1} \otimes I_\alpha](f) \\
&= (\rho'_{1,1}F_1 \otimes I_\alpha)(f) && \text{by (\#)} \\
&= (\rho'_{1,1} \otimes I_\alpha)(F_1 \otimes I_\alpha)(f) \\
&= (\rho'_{1,1} \otimes I_\alpha)\overline{F}_\alpha(f)
\end{aligned}$$

Now we shall prove that $\overline{F}$ is $\pi-$comodule map, which follows from, for $f \in Ind(\rho)_{\alpha\beta}$,

$$\begin{aligned}
&(\overline{F}_\alpha \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})(f) \\
&= (F_1 \otimes I_\alpha \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})(f) \\
&= (I_1 \otimes \Delta_{\alpha,\beta})\overline{F}_\alpha(f).
\end{aligned}$$

∎

Corollary 5.2. *If (V,ρ) and (W,ρ') are equivalent right $\pi-$comodules over C, then the induced representations $Ind(\rho)$ and $Ind(\rho')$ are equivalent right $\pi-$comodules over H.*

Lemma 5.3. *Let (V,ρ) and (W,ψ) be two right $\pi-$comodules over C, then $V\oplus W=\{(V\oplus W)_\alpha=V_\alpha\oplus W_\alpha\}_{\alpha\in\pi}$ is right $\pi-$comodule over C by $\rho\oplus\psi=\{(\rho\oplus\psi)_{\alpha,\beta}=\rho_{\alpha,\beta}\oplus\psi_{\alpha,\beta}:V_{\alpha\beta}\oplus W_{\alpha\beta}\to(V_\alpha\oplus W_\alpha)\otimes C_\alpha\}_{\alpha,\beta\in\pi}$ where*

$$\rho_{\alpha,\beta}\oplus\psi_{\alpha,\beta}=(l_\alpha^V\otimes I_\beta)\rho_{\alpha,\beta}P_{\alpha\beta}^V+(l_\alpha^W\otimes I_\beta)\psi_{\alpha,\beta}P_{\alpha\beta}^W,\ \ \textit{where}$$

$$P_\alpha^V:V_\alpha\oplus W_\alpha\to V_\alpha$$

$$P_\alpha^W:V_\alpha\oplus W_\alpha\to W_\alpha$$

are the natural projection maps and

$$l_\alpha^V:V_\alpha\to V_\alpha\oplus W_\alpha$$

$$l_\alpha^W:W_\alpha\to V_\alpha\oplus W_\alpha$$

are the natural immersion maps.

Theorem 5.4. *Let (V,ρ) and (W,ψ) be two right $\pi-$comodules over C, then*

$$Ind(\rho\oplus\psi)\cong Ind(\rho)\oplus Ind(\psi).$$

Proof. We define $\theta:Ind(\rho\oplus\psi)\to Ind(\rho)\oplus Ind(\psi)$ as follows: $\theta=\{\theta_\alpha=(P_1^V\otimes I_\alpha)+(P_1^W\otimes I_\alpha)\}_{\alpha\in\pi}$. θ_α is an isomorphism of vector spaces and $\theta_\alpha:Ind(\rho\oplus\psi)_\alpha\to Ind(\rho)_\alpha\oplus Ind(\psi)_\alpha$. We will prove that, for $f\in Ind(\rho\oplus\psi)_\alpha$,

1. $(P_1^V\otimes I_\alpha)(f)\in Ind(\rho)_\alpha$
2. $(P_1^W\otimes I_\alpha)(f)\in Ind(\psi)_\alpha$

and hence $\theta_\alpha(f)\in Ind(\rho)_\alpha\oplus Ind(\psi)_\alpha$.

$$\begin{aligned}
&(I_1\otimes L_{1,\alpha})(P_1^V\otimes I_\alpha)(f)\\
&=(P_1^V\otimes I_1\otimes I_\alpha)(I_1\otimes L_{1,\alpha})(f)\\
&=(P_1^V\otimes I_1\otimes I_\alpha)((\rho\oplus\psi)_{1,1}\otimes I_\alpha)(f)\\
&=(P_1^V\otimes I_1\otimes I_\alpha)[[(l_\alpha^V\otimes I)\rho_{1,1}P_1^V+(l_1^W\otimes I)\psi_{1,1}P_1^W]\otimes I_\alpha](f)\\
&=(P_1^V\otimes I_1\otimes I_\alpha)[(l_\alpha^V\otimes I)\rho_{1,1}P_1^V\otimes I_\alpha](f)+(P_1^V\otimes I_1\otimes I_\alpha)[(l_\alpha^W\otimes I)\psi_{1,1}P_1^W\otimes I_\alpha](f)\\
&=(P_1^V\otimes I_1\otimes I_\alpha)[(l_\alpha^V\otimes I)\rho_{1,1}P_1^V\otimes I_\alpha](f)\\
&=[(P_1^Vl_\alpha^V\otimes I)\rho_{1,1}P_1^V\otimes I_\alpha](f)\\
&=(\rho_{1,1}P_1^V\otimes I_\alpha)(f)\\
&=(\rho_{1,1}\otimes I_\alpha)(P_1^V\otimes I_\alpha)(f)
\end{aligned}$$

then $(P_1^V \otimes I_\alpha)(f) \in Ind(\rho)_\alpha$.

Similar $(P_1^W \otimes I_\alpha)(f) \in Ind(\psi)_\alpha$. Therefore $\theta_\alpha(f) \in Ind(\rho)_\alpha \oplus Ind(\psi)_\alpha$. Clear θ_α is an isomorphism. The intertwining property is the commutativity of the following diagram

$$\begin{tikzcd}
Ind(\rho \oplus \psi)_{\alpha\beta} \arrow[r, "\theta_{\alpha\beta}"] \arrow[d, "I \otimes \Delta_{\alpha,\beta}"'] & Ind(\rho)_{\alpha\beta} \oplus Ind(\psi)_{\alpha\beta} \arrow[d, "\overline{\rho}_{\alpha,\beta} \oplus \overline{\psi}_{\alpha,\beta}"] \\
Ind(\rho \oplus \psi)_\alpha \otimes H_\beta \arrow[r, "\theta_\alpha \otimes I_\beta"'] & (Ind(\rho)_\alpha \oplus Ind(\psi)_\alpha) \otimes H_\beta
\end{tikzcd}$$

where $Ind(\rho \oplus \psi)$ is right $\pi-$comodule over H by $I \otimes \Delta = \{I_{V_1 \oplus W_1} \otimes \Delta_{\alpha,\beta}\}$. Also, $Ind(\rho) \oplus Ind(\psi)$ is right $\pi-$comodule over H by

$$\overline{\rho} \oplus \overline{\psi} = \{\overline{\rho}_{\alpha,\beta} \oplus \overline{\psi}_{\alpha,\beta}\}$$

where

$$\overline{\rho}_{\alpha,\beta} \oplus \overline{\psi}_{\alpha,\beta} = (l_\alpha^{Ind(\rho)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\rho)} + (l_\alpha^{Ind(\psi)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\psi)}$$

Now,

$$\begin{aligned}
&[\overline{\rho}_{\alpha,\beta} \oplus \overline{\psi}_{\alpha,\beta}]\vartheta_{\alpha\beta}(f) \\
&= [(l_\alpha^{Ind(\rho)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\rho)} + (l_\alpha^{Ind(\psi)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\psi)}] \\
&\quad [(P_1^V \otimes I_{\alpha\beta}) + (P_1^W \otimes I_{\alpha\beta})][(v + w) \otimes h] \\
&= [(l_\alpha^{Ind(\rho)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\rho)} + (l_\alpha^{Ind(\psi)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\psi)}][(v \otimes h) + (w \otimes h)] \\
&= [(l_\alpha^{Ind(\rho)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\rho)}][(v \otimes h) + (w \otimes h)] + [(l_\alpha^{Ind(\psi)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})P_{\alpha\beta}^{Ind(\psi)}] \\
&\quad [(v \otimes h) + (w \otimes h)] \\
&= [(l_\alpha^{Ind(\rho)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})](v \otimes h) + [(l_\alpha^{Ind(\psi)} \otimes I_\beta)(I_1 \otimes \Delta_{\alpha,\beta})](w \otimes h) \\
&= (v \otimes h_1 + 0) \otimes h_2 + (0 + w \otimes h_1) \otimes h_2 \\
&= [(v \otimes h_1) + (w \otimes h_1)] \otimes h_2 \\
&= [[(P_1^V \otimes I_\alpha) + (P_1^W \otimes I_\alpha)] \otimes I_\beta][(v + w) \otimes \Delta_{\alpha,\beta}(h)] \\
&= (\theta_\alpha \otimes I_\beta)(I \otimes \Delta_{\alpha,\beta})(f)
\end{aligned}$$

i.e., the above diagram commutes.

▮

Corollary 5.5. *Let H be a Hopf $\pi-$coalgebra, (C, σ) a $\pi-$coisotropic quantum subgroup of H and (V, ρ) a right $\pi-$comodules over C. If $Ind(\rho)$ is simple, then ρ is simple.*

The inverse of the above result is not true in general case by the following example.

Example Let A be the algebra generated by 1 and the elements x, y, x^{-1}, y^{-1} and g with

$$\begin{aligned}&xx^{-1} = x^{-1}x \; = \; yy^{-1} = y^{-1}y = 1, \quad xy = yx,\\ &\quad xg = qgx, \quad yg = qgy, \quad x^{-1}g = q^{-1}gx^{-1}, \quad y^{-1}g = q^{-1}gy^{-1}\end{aligned}$$

for some fixed non zero $q \in K$, then A become Hopf algebra with

$$\Delta \begin{pmatrix} g \\ x^{\pm 1} \\ y^{\pm 1} \end{pmatrix} = \begin{pmatrix} g \otimes x + y \otimes g \\ x^{\pm 1} \otimes x^{\pm 1} \\ y^{\pm 1} \otimes y^{\pm 1} \end{pmatrix} \quad , \quad \epsilon \begin{pmatrix} g \\ x^{\pm 1} \\ y^{\pm 1} \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix},$$

$$S \begin{pmatrix} g \\ x^{\pm 1} \\ y^{\pm 1} \end{pmatrix} = \begin{pmatrix} -y^{-1}gx^{-1} \\ x^{\mp 1} \\ y^{\mp 1} \end{pmatrix}$$

Now, we take π the set of all group like elements of A, then π acts on A by Hopf algebra endomorphism as $\lambda : \pi \to End(A)$ where $\lambda(\alpha)(\beta) = \alpha\beta\alpha^{-1}$ for all $\alpha, \beta \in \pi$. Then, by [T], we can construct Hopf $\pi-$coalgebra as $H = \{H_\alpha\}_{\alpha\in\pi}$ where for each $\alpha \in \pi$, the algebra H_α is a copy of A with comultiplication $\Delta_{\alpha,\beta}(i_{\alpha\beta}(a)) = i_\alpha(\lambda(\beta)(a_1)) \otimes i_\beta(a_2)$, counit $\epsilon_1(i_1(a)) = \epsilon(a)$ and the antipode $S_\alpha(i_\alpha(a)) = i_{\alpha^{-1}}(\lambda(\alpha)(s(a)))$ where $a \in A$, $\Delta(a) = a_1 \otimes a_2$ is the given comultiplication in A written in Sweedler‘s sigma notation.

Now take $J = \{J_\alpha\}_{\alpha\in\pi}$ where for each $\alpha \in \pi$, J_α is a copy of J_R where J_R is a right ideal of A generated by $x - 1, y - 1, x^{-1} - 1$ and $y^{-1} - 1$. J_R is a coideal becouse, for example,

$$\begin{aligned}&\Delta(x-1) = x \otimes x - 1 \otimes 1 = (x-1) \otimes x + 1 \otimes (x-1)\\ &\;\epsilon(x-1) = \epsilon(x) - \epsilon(1) = 0 - 0 = 0,\end{aligned}$$

J is a family of right ideals and also J is $\pi-$ coideal. Then, by theorem 1.2, $C = H/J$ is $\pi-$ coalgebra such that $\overline{\sigma} = \{\overline{\sigma}_\alpha = \sigma : H_\alpha \to C_\alpha = H_\alpha/J_\alpha\}_{\alpha\in\pi}$ is $\pi-$coalgebra epimorphism map. Furthermore, every C_α is $H_\alpha-$ module such that every $\overline{\sigma}_\alpha$ is module map.

Now take $V = \{V_\alpha\}_{\alpha\in\pi}$ is simple $\pi-$comodule over C by $\rho = \{\rho_{\alpha,\beta}\}_{\alpha,\beta\in\pi}$, this implies that V_1 is simple C_1-comodule by $\rho_{1,1}$. Since C_1 is cocommutative, then V_1 has one dimensional. So, we can suppose that $\rho_{1,1}(w) = w \otimes 1$. The induced representation in this case is $Ind(\rho) = \{Ind(\rho)_\alpha\}$ where

$Ind(\rho)_\alpha$ is generated by x, x^{-1}, y and y^{-1}. Also, $Ind(\rho)_\alpha \subseteq H_\alpha$. Now, take $B = \{B_\alpha\}_{\alpha\in\pi}$ where B_α is a subalgebra of H_α generated by x, and x^{-1}. Clear B is subHopf $\pi-$coalgebra of H and every $B_\alpha \subseteq Ind(\rho)_\alpha$ and every element in B_α is group like, the B is non trivial $\pi-$ subcomodule of $Ind(\rho)$ and hence $Ind(\rho)$ is not simple $\pi-$comodule over H.

Now, we return to study the equivalent and direct sum of coinduced representation.

Theorem 5.6. *Let (V,ρ) and (W,ρ') be a right $\pi-$comodules over a $\pi-$coisotropic quantum subgroup (C,σ). If there exist a vector space isomorphism $F_1 : V_1 \to W_1$ such that*

$$\psi_{1,1}F_1 = (F_1 \otimes I_1)\rho_{1,1} \qquad \text{(1)}$$

then the coinduced representations $Coind(\rho)$ and $Coind(\psi)$ are equivalent right $\pi-$comodules over H.

Proof. Let us consider the vector space isomorphisms $\overline{F} = \{\overline{F}_\alpha : Coind(\psi)_\alpha \to Coind(\rho)_\alpha\}_{\alpha\in\pi}$, where

$$\overline{F}_\alpha(f) = f \circ F_1 \quad \text{for all} \quad f \in Coind(\psi)_\alpha.$$

Firstly, we must be prove $\overline{F}_\alpha(f) = f \circ F_1 \in Coind(\rho)_\alpha$, for $\alpha \in \pi$ for $\alpha \in \pi$, $f \in Coind(\psi)_\alpha$.

$$\begin{aligned}
& L_{1,\alpha}\overline{F}_\alpha(f) \\
&= L_{1,\alpha} f \circ F_1 \\
&= (I \otimes f)\psi_{1,1}F_1 \\
&= (I \otimes f)(I \otimes F_1)\rho_{1,1} && \text{by (1)} \\
&= (I \otimes f \circ F_1)\rho_{1,1} \\
&= (I \otimes \overline{F}_\alpha(f))\rho_{1,1}.
\end{aligned}$$

Now, we will prove the following diagram commutes

$$\begin{tikzcd}
Coind(\rho)_{\alpha\beta} \arrow[r, "\overline{F}_{\alpha\beta}"] \arrow[d, "\Omega_{\alpha,\beta}"'] & Coind(\rho)_{\alpha\beta} \arrow[d, "\overline{\Omega}_{\alpha,\beta}"] \\
Coind(\psi)_\alpha \otimes H_\beta \arrow[r, "\overline{F}_\alpha \otimes I_\beta"'] & Coind(\rho)_\alpha \otimes H_\beta
\end{tikzcd}$$

For $h \in Coind(\psi)_{\alpha\beta}$

$$\begin{aligned}
&\overline{\Omega}_{\alpha,\beta}\overline{F}_{\alpha\beta}(h)\\
&= \overline{\Omega}_{\alpha,\beta}(h \circ F_1)\\
&= \sim (I_\alpha \otimes g^\beta)\Delta_{\alpha,\beta}(h \circ F_1) \otimes e^\beta\\
&= \overline{\Omega}_\alpha(\sim (I_\alpha \otimes g^\beta)\Delta_{\alpha,\beta}h) \otimes e^\beta\\
&= (\overline{\Omega}_\alpha \otimes I_\beta)((\sim (I_\alpha \otimes g^\beta)\Delta_{\alpha,\beta}h \otimes e^\beta)\\
&= (\overline{\Omega}_\alpha \otimes I_\beta)\Omega_{\alpha,\beta}(h).
\end{aligned}$$

Now we will prove $\overline{F}_\alpha$ is an isomorphism. Let $t \in Coind(\rho)_\alpha$, there exist $t \circ F_1^{-1} \in Coind(\psi)_\alpha$ such that $\overline{F}_\alpha(t \circ \overline{F}_1^{-1}) = t\overline{F}_1^{-1}F_1 = t$. Therefore $\overline{F}_\alpha$ is onto, by finite dimensional $\overline{F}_\alpha$ is an isomorphism.

The remaining is prove $t \circ F_1^{-1} \in Coind(\psi)_\alpha$, i.e., $L_{1,\alpha}t \circ F_1^{-1} = (I \otimes t \circ F_1^{-1})\psi_{1,1}$ i.e., $(I \otimes t)\rho_{1,1}F_1^{-1} = (I \otimes t \circ F_1^{-1})\psi_{1,1}$. We have $\psi_{1,1}F_1 = (I \otimes F_1)\rho_{1,1}$ implies $(I \otimes F_1^{-1})\psi_{1,1}F_1 = \rho_{1,1}$ that is means

$$(I \otimes F_1^{-1})\psi_{1,1}F_1 = \rho_{1,1}F_1^{-1}F_1$$

and hence

$$[(I \otimes F_1^{-1})\psi_{1,1} - \rho_{1,1}F_1^{-1}]F_1 = 0$$

implies that

$$[(I \otimes F_1^{-1})\psi_{1,1} - \rho_{1,1}F_1^{-1}]F_1(v) = 0 \qquad \text{for all} \quad v \in V_1.$$

Since F_1 is onto, then

$$[(I \otimes F_1^{-1})\psi_{1,1} - \rho_{1,1}F_1^{-1}](w) = 0 \quad \text{for all} \quad w \in W_1,$$

that is $(I \otimes F_1^{-1})\psi_{1,1} - \rho_{1,1}F_1^{-1} = 0$ i.e., $(I \otimes F_1^{-1})\psi_{1,1} = \rho_{1,1}F_1^{-1}$ therefore

$$(I \otimes t)(I \otimes F_1^{-1})\psi_{1,1} = (I \otimes t)\rho_{1,1}F_1^{-1},$$

then

$$(I \otimes t \circ F_1^{-1})\psi_{1,1} = L_{1,\alpha}t \circ F_1^{-1}.$$

Thus $Coind(\psi)$ and $Coind(\rho)$ are equivalent. ∎

Corollary 5.7. *If (V, ρ) and (W, ψ) are equivalent right $\pi-$comodules over C, then the coinduced representations $Coind(\rho)$ and $Coind(\psi)$ are equivalent right $\pi-$comodules over H.*

Theorem 5.8. *Suppose (V, ρ) and (W, ψ) are left $\pi-$comodules over C, then*

$$Coind(\rho \oplus \psi) \cong Coind(\rho) \oplus Coind(\psi).$$

Proof. Since (V, ρ) and (W, ψ) are left $\pi-$comodules over C, then $(Coind(\rho), \Omega)$ and $(Coind(\psi), \Upsilon)$ are right $\pi-$comodules over H and hence, by lemma 5.3, $(Coind(\rho) \oplus (Coind(\psi))$ is right $\pi-$comodule over H by $\Lambda = \{\Lambda_{\alpha,\beta}\}$ where

$$\Lambda_{\alpha,\beta} = (l_\alpha^{Coind(\rho)} \otimes I_\beta)\Omega_{\alpha,\beta} P_{\alpha\beta}^{Coind(\rho)} + (l_\alpha^{Coind(\psi)} \otimes I_\beta)\Upsilon_{\alpha,\beta} P_{\alpha\beta}^{Coind(\psi)}.$$

Also, by lemma 5.3, $V \oplus W = \{V_\alpha \oplus W_\beta\}$ is left $\pi-$comodule over C by

$$\rho \oplus \psi = \{(\rho \oplus \psi)_{\alpha,\beta} = \rho_{\alpha,\beta} \oplus \psi_{\alpha,\beta} : V_{\alpha\beta} \oplus W_{\alpha\beta} \to C_\alpha \otimes (V_\beta \oplus W_\beta)\}_{\alpha,\beta\in\pi}$$

where

$$\rho_{\alpha,\beta} \oplus \psi_{\alpha,\beta} = (I_\alpha \otimes l_\beta^V)\rho_{\alpha,\beta} P_{\alpha\beta}^V + (I_\alpha \otimes l_\beta^W)\psi_{\alpha,\beta} P_{\alpha\beta}^W.$$

Then $Coind(\rho \oplus \psi)$ is right $\pi-$comodule over H by $\Gamma = \{\Gamma_{\alpha,\beta}\}$.

Now, we define $\chi = \{\chi_\alpha : Coind(\rho \oplus \psi)_\alpha \to Coind(\rho)_\alpha \oplus Coind(\psi)_\alpha\}_{\alpha\in\pi}$, where

$$\chi_\alpha(g) = g \circ l_1^V + g \circ l_1^W \quad \text{for all} \quad g \in Coind(\rho \oplus \psi)_\alpha.$$

Firstly, we will prove $g \circ l_1^V \in Coind(\rho)_\alpha$

$$\begin{aligned}
&L_{1,\alpha} g \circ l_1^V \\
&= (L_{1,\alpha} g) l_1^V \\
&= (I_1 \otimes g)(\rho_{1,1} \otimes \psi_{1,1}) l_1^V \\
&= (I_1 \otimes g)[(I_1 \otimes l_1^V)\rho_{1,1} P_1^V + (I_1 \otimes l_1^W)\psi_{1,1} P_1^W] l_1^V \\
&= (I_1 \otimes g)(I_1 \otimes l_1^V)\rho_{1,1} P_1^V l_1^V \\
&= (I_1 \otimes g)(I_1 \otimes l_1^V)\rho_{1,1} \\
&= (I_1 \otimes g \circ l_1^V)\rho_{1,1}.
\end{aligned}$$

Similar $g \circ l_1^W \in Coind(\psi)_\alpha$, and hence $\chi_\alpha(g) \in Coind(\rho)_\alpha \oplus Coind(\psi)_\alpha$.

We will prove χ_α is 1-1 (and hence isomorphism by finite dimensional). Suppose $\chi_\alpha(g) = 0$, then $(\chi_\alpha(g))(v_1 + w_1) = 0$ for all $v_1 \in V_1$ and $w_1 \in W_1$ implies $g(v_1 + w_1) = 0$ for all $v_1 \in V_1$ and $w_1 \in W_1$, then $g = 0$.

Finally, we will prove that the following diagram commutes

$$\begin{tikzcd}
Coind(\rho \oplus \psi)_{\alpha\beta} \arrow[r, "\chi_{\alpha,\beta}"] \arrow[d, "\Gamma_{\alpha,\beta}"'] & Coind(\rho)_{\alpha\beta} \oplus Coind(\psi)_{\alpha\beta} \arrow[d, "\Lambda_{\alpha,\beta}"] \\
Coind(\rho \oplus \psi)_\alpha \otimes H_\beta \arrow[r, "\chi_\alpha \otimes I_\beta"'] & Coind(\rho)_\alpha \oplus Coind(\psi)_\alpha \otimes H_\beta
\end{tikzcd}$$

$$
\begin{aligned}
&(\chi_\alpha \otimes I_\beta)\Gamma_{\alpha,\beta}(g)\\
&= (\chi_\alpha \otimes I_\beta)[\sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta} g \otimes e^\beta]\\
&= [(\sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta} g) l_1^V + (\sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta} g) l_1^W] \otimes e^\beta\\
&= [\sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta}(g \circ l_1^V + g \circ l_1^W)] \otimes e^\beta
\end{aligned}
$$

On other hand

$$
\begin{aligned}
&\Lambda_{\alpha,\beta}\chi_{\alpha,\beta}(g)\\
&= [(l_\alpha^{Coind(\rho)} \otimes I_\beta)\Omega_{\alpha,\beta} P_{\alpha\beta}^{Coind(\rho)} + (l_\alpha^{Coind(\psi)} \otimes I_\beta)\Upsilon_{\alpha,\beta} P_{\alpha\beta}^{Coind(\psi)}](g l_1^V + g l_1^W)\\
&= (l_\alpha^{Coind(\rho)} \otimes I_\beta)\Omega_{\alpha,\beta} g \circ l_1^V + (l_\alpha^{Coind(\psi)} \otimes I_\beta)\Upsilon_{\alpha,\beta} g \circ l_1^W\\
&= (l_\alpha^{Coind(\rho)} \otimes I_\beta)[\sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta} g \circ l_1^V \otimes e^\beta] + (l_\alpha^{Coind(\psi)} \otimes I_\beta)[\sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta} g \circ l_1^W \otimes e^\beta]\\
&= [\sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta} g \circ l_1^V \oplus 0] \otimes e^\beta + [0 \oplus \sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta} g \circ l_1^W] \otimes e^\beta\\
&= \sim (I_\alpha \otimes f^\beta)\Delta_{\alpha,\beta}(g \circ l_1^V + g \circ l_1^W) \otimes e^\beta.
\end{aligned}
$$

▮

Corollary 5.9. *Let H be a Hopf $\pi-$coalgebra, (C, σ) a $\pi-$coisotropic quantum subgroup of H and (V, ρ) a right $\pi-$comodules over C. If $Coind(\rho)$ is simple, then ρ is simple.*

References

[C] N. Ciccoli, Induction of quantum group representation, J. Geometry and Physics, (1-16), 1998.

[G-I] A.Gonzalez-Ruiz, L. A. Ibort, Induction of Quantum Groups, Phys. Lett. B., (296) (107-110), 1992.

[H] A. S. Hegazi, M. G. El-Agawany, Fatma Ismail and Ibrahim Saleh, On the representation theory of quantum groups, M. Sc. Thesis Cairo Uni. 2001.

[M] S. Majid, Foundation of quantum group theory, Cambridge Univeristy Press, 1995.

[Mon] S. Montgomery, Hopf algebras and their actions on rings, Amer. Math. Soc., 1993.

[S] M. E. Sweedler, Hopf algebras, Cornell Uni. New York,1969.

[T] V. G. Truaev, Homotopy field theory in dimension 3 and crossed group-categories, preprent GT/0005291.

[V] A. Virelizier, Hopf group-coalgebras, J. of Pure and Applied Algebra, 171 (75-122), 2002.

[V1] A. Virelizier, Involutory Hopf group-coalgebras and flat bundles over 3 manifolds, Arxiv:Math. GT/0206254 v1 24 Jun. 2002.